

# Centennial Valley Arctic Grayling Adaptive Management Project Annual Report, 2018

12 April, 2019




## EXECUTIVE SUMMARY

Arctic grayling (*Thymallus arcticus*) are a freshwater holarctic salmonid that were once widespread throughout the Upper Missouri River (UMR) drainage as a glacial relict population. One of the last endemic grayling populations remaining in the UMR drainage resides in the Centennial Valley (CV) of southwestern Montana. Spawning is largely limited to Red Rock, Corral, Elk Springs, and Odell creeks, with Red Rock Creek likely supporting 80-90% of annual spawning in the CV. It is presumed that most of the grayling population in the CV spends non-breeding portions of the year in Upper Red Rock Lake (Upper Lake). Red Rock Lakes National Wildlife Refuge (Refuge) encompasses Upper Lake, and nearly all of the currently occupied grayling spawning habitat within Red Rock, Elk Springs, and Odell creeks.

The estimated number of grayling in the 2018 Red Rock Creek spawning population was 387 (95% CI = 168–682), nearly unchanged from the previous year ( $\hat{N} = 176$ , 95% CI = 159–213; Figure 1). Suitable habitat the prior winter within Upper Lake (i.e., water depth below the

ice  $\geq 1$  m and dissolved oxygen  $\geq 4$  ppm) reached a minimum during February sampling at an estimated 19 ha. This falls within a range of suitable winter habitat (i.e., 10–25 ha) previously hypothesized to lead to reduced grayling survival as documented in 2016. Area of winter habitat for the current year (2019) was estimated in January as 0 ha; 1995 was the only other year 0 ha of suitable winter habitat was estimated.

Suitable spawning habitat was quantified in 2017, with an estimated total area of suitable spawning habitat ( $A_{ts}$ ) of 0 ha, and weighted area of suitable habitat ( $A_{tw}$ ) of 4 ha, in Red Rock and Elk Springs creeks (see **METHODS** below for description of variables). Barriers precluding grayling from accessing suitable spawning habitat, i.e., beaver dams, were notched prior to spawning again in 2018 (similar to 2017) as part of the second hypothesis test of the Adaptive Management Plan. Hydrology during the 2018 breeding season critical period (i.e., *peak spawning + 188 degree days + 5 weeks*) was characterized using four variables related to stream flows (cubic feet second<sup>-1</sup> [cfs]) and temperature (°C) – mean daily discharge ( $mdd = 111.4$  cfs), cumulative degree days from peak fry emergence to five weeks post-emergence ( $cdd = 364.2$ ), days above bankfull discharge ( $cbf = 15$ ), and days above 67% of bankfull discharge ( $c67bf = 38$ ). These values represent 133%, 89%, 213%, and 186% of the long-term mean values for  $mdd$ ,  $cdd$ ,  $cbf$ , and  $c67bf$ , respectively.



**Figure 1.** Estimated annual Red Rock Creek Arctic grayling spawning population abundance and 95% confidence intervals (when available), 1994–2018. Confidence intervals are not available for some years prior to initiation of the Adaptive Management Plan.

The first management experiment conducted as part of the Adaptive Management Plan was reducing non-native Yellowstone cutthroat trout (*Oncorhynchus clarkii bowieri*). Trout were

captured and euthanized at a fish weir and fishing regulations were liberalized to 20 cutthroat trout per day (excluding a stream closure 15 May–14 June) on Red Rock Creek. Both actions were first implemented in 2013; the weir was operated through 2016 and liberalized harvest occurred through 2017. During 2013–2017 a total of 7149 cutthroat trout were removed from Red Rock Creek. Removal efforts peaked in 2014 at 2604 cutthroat removed. The first year a grayling response to cutthroat removal could be quantified was 2016 when the 2013 grayling cohort recruited.

The second management experiment is exploring the relationship between spawning habitat and grayling recruitment. Efforts beginning in 2017 increased suitable habitat available for spawning by 1) ensuring fish passage at beaver dams ( $n > 50$  in 2016) via notching, 2) restoration of an Elk Springs Creek channel that will provide direct connection to Upper Lake, and 3) restoration of spawning habitat at the head of Elk Springs Creek. This experiment will be undertaken for 4–5 years, similar to the cutthroat trout reduction. Increased per capita availability of spawning habitat is hypothesized to increase egg ( $\beta$ ) and age-0 fish in-stream ( $\gamma$ ) survival; the first year a possible response could be quantified is 2020, when the 2017 grayling cohort recruits.

Learning in the context of this project occurs through comparison of model predictions with reality (i.e., predicted grayling population vs. actual grayling population). Each hypothesized driver of grayling population dynamics, i.e., winter habitat, spawning habitat, non-native fish, and spring hydrology, is represented by a model structured to estimate the driver’s influence on a specific life stage. Grayling recruit at age-3, which necessitates having three consecutive years of data to make model-based predictions. The first time-series of data necessary to make a prediction was available in 2016, which was also the first population estimate that could be used to fit models. However, it is preferable to fit models using several population estimates. We addressed this issue in the short-term using simulations to estimate the influence of each hypothesized driver of grayling population (see **Methods**). Based on simulation results, the *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models predicted 67, 92, and 840 grayling, respectively, in the 2018 Red Rock Creek spawning population. Model weights, i.e., relative support for a model given the data, were 0.587, 0.311, 0.103 for the *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models, respectively.

---

## Authors:

Jeff Warren<sup>1</sup>, Matt Jaeger<sup>2</sup>, Andrew Gilham<sup>3</sup>, Kyle Cutting<sup>4</sup>, Lucas Bateman<sup>2</sup>, Terrill Paterson<sup>2</sup>, and Mike Duncan<sup>2</sup>.

<sup>1</sup> U.S. Fish and Wildlife Service, Division of Science Resources,

<sup>2</sup> Montana Fish, Wildlife and Parks,

<sup>3</sup> U.S. Fish and Wildlife Service, Montana Fish and Wildlife Conservation Office,

<sup>4</sup> U.S. Fish and Wildlife Service, Red Rock Lakes National Wildlife Refuge.

# INTRODUCTION

Montana Arctic grayling (*Thymallus arcticus*; hereafter grayling) were patchily distributed throughout the Upper Missouri River (UMR) drainage prior to the mid-1850s. This population declined to about 4% of their perceived historic distribution by the 1990s, which led to formal consideration for listing under the Endangered Species Act (USFWS 2014). One of the last populations of indigenous UMR grayling resides in the Centennial Valley (CV) of southwestern Montana. Grayling were historically distributed among at least a dozen CV streams and three lakes at presumably high abundances (Nelson 1954). Perceived distribution and abundance declined to historic lows sometime between the 1950s and mid-1990s, but have since improved, although large fluctuations in abundance still occur (USFWS 2014, MAGWG in press).

Currently, most of the CV grayling population spawns in Red Rock Creek and spends non-breeding portions of the year in Upper Red Rock Lake (Upper Lake) within Red Rock Lakes National Wildlife Refuge (Refuge; Figure 2). Over the past 70 years numerous hypotheses were posited regarding drivers of the CV grayling population, including 1) reduction and alteration of spawning habitat, 2) predation by, and competition with, non-native fishes, and 3) limited winter habitat. Although these hypotheses have been repeatedly proposed to explain population fluctuations, drivers of the population remain unclear.


Previous and ongoing research has focused on aspects of each hypothesis but has not linked them to demographic responses in grayling, which precludes inference regarding their role as population drivers. Resultantly, the most effective management and conservation approaches for CV grayling remain ambiguous, and selecting management actions can be contentious among and within agencies. This plan seeks to elucidate the relative effect of hypothesized drivers of CV grayling abundance to direct future management of this population. Determining the cause of previous population declines, *per se*, is not the primary issue of grayling conservation and management – finding an effective strategy to achieve population goals and prevent future declines is. In an effort to accomplish this, an adaptive management (AM) approach is being undertaken (Walters 1986). The Centennial Valley Arctic Grayling Adaptive Management Plan (AMP) embraces existing uncertainty regarding drivers of the grayling population in the CV to provide further understanding of important limiting factors and help guide management actions toward those that will have the most direct benefit to grayling.

Due to the initial level of structural uncertainty, and agency conflict regarding that uncertainty, the AMP was divided into two phases – a ‘management as experiment’ phase that emphasizes learning, i.e., reducing structural uncertainty (MacNab 1983, Walters 1986, Walters and Holling 1990), and an active adaptive management phase. The former was designed to explore grayling population response to hypothesized drivers that could be influenced via management actions. The latter will use the information gained in phase 1 to determine an optimal policy to inform annual management decisions (while still learning, but with less of an emphasis on learning).

## STUDY AREA

The Centennial Valley of southwestern Montana is a high-elevation (ca. 2013 m) valley dominated by sagebrush steppe comprising *Artemisia* spp. shrub overstory and native bunchgrass understory (e.g., *Festuca* spp., *Nasella* spp., and *Hesperostipa* spp.). The valley is bounded on the north by the north-south trending Gravelly and Snowcrest mountain ranges and on the south by the east-west trending Centennial Mountains. Extensive wetlands exist throughout the CV, including a large shallow lake/wetland complex encompassed by Red Rock Lakes National Wildlife Refuge (Figure 2). The complex comprises Upper Red Rock, Lower Red Rock, and Swan lakes and associated palustrine emergent marsh dominated by seasonally-flooded sedge (*Carex* spp.). The complex is a remnant of Pleistocene Lake Centennial, a prehistoric lake that was believed to have formerly covered the valley floor to a depth of ca. 20 m (Mumma 2010). Upper Lake, the largest and deepest of the lakes, is ca. 1198 ha with a maximum depth of 2 m. The geologic (Sonderegger 1981; Centennial Valley Historical Society 2006), hydrologic (Deeds and White 1926, MTFWP 1989, MCA 2000), and fisheries (Nelson 1952, Randall 1978, Boltz 2000, Oswald et al. 2008) resources and contemporary administrative status (USFWS 2009) within the Centennial Valley are well described elsewhere.

The Centennial Valley includes all tributaries of the Red Rock River and their associated drainages upstream of Lima Dam (Figure 2). Most Upper Lake tributaries have their origins to the south along the eastern extent of the Centennial Mountains. Red Rock Creek, the largest of these tributaries, originates at an elevation of 2,562 m and flows north and west ca. 21 km to the northeast shore of Upper Lake. Elk Springs Creek originates from a series of springs south of Elk Lake and flows southwest, entering Upper Lake along the northeast shore. Red Rock River exits Upper Lake in the northwest corner, carrying water through the River Marsh and into the northwestern corner of Lower Lake. Red Rock River continues westward through the outlet of Lower Lake, ca. 1.5 km west of where it enters the lake, leaving the CV near Lima, MT after passing through the 13 km long Lima Reservoir. Long Creek enters the Red Rock River 17 km downstream of the Lower Lake outlet and just upstream of Lima Reservoir.



**Figure 2.** Arctic grayling Adaptive Management Plan study area within the Centennial Valley of southwestern Montana.

## METHODS

### Grayling Abundance and Survival

Grayling and cutthroat trout were captured by mobile anode electrofishing (May and June) and a stationary weir (April–June) from 2013 to 2017 (see Paterson 2013 for further description). Beginning in 2018 only electrofishing surveys occurred. Grayling were uniquely marked with a visual implant (VI) tag, sexed, and length ( $\pm 1$  mm) and weight ( $\pm 1$  g) recorded. We used capture-mark-recapture (CMR) models implemented in program MARK (White and Burnham 1999) using the RMark package (Laake 2013) in R version 3.5.2 (R Development Core Team 2018) to estimate grayling abundance with closed population models and apparent survival ( $\phi$ ) with open population models. Apparent survival confounds permanent emigration and mortality; therefore survival will be biased low if adult grayling fidelity to spawning streams is  $< 1$ .

### Models of System Dynamics

Spawning habitat, non-native fishes, winter habitat, and spring and summer hydrology have all been identified as potentially important drivers of grayling population dynamics in the

upper CV. Each of these hypotheses is translated into a model, or set of models, to link hypothesized drivers and demographic rates.

The annual abundance of spawning grayling is the product of demographic rates ranging from adult survival to the number of eggs deposited per female (fecundity) three years prior. All population models for spawning grayling share a common balance equation that allows prediction of annual abundance as a function of survival and recruitment processes:

$$N_{t+1} = N_t S_t + (F_{t-2} \alpha_{t-2} \beta_{t-2} \gamma_{t-2} \delta_{t-2} \epsilon_{t-1}) \theta_t \quad (1)$$

The number and survival of adult (i.e., reproductive age) grayling in year  $t$  is  $N_t$  and  $S_t$ , respectively. Assuming recruitment occurs with the age-3 cohort in year  $t + 1$  (i.e., knife-edge recruitment at age-3), the number of potential age-2 recruits in year  $t$  is the product of:

- $F_{t-2}$  – the number of females in the spawning run in year  $t - 2$ ,
- $\alpha_{t-2}$  – length-specific fecundity rate, year  $t - 2$ ,
- $\beta_{t-2}$  – probability of an egg being fertilized and hatching, year  $t - 2$ ,
- $\gamma_{t-2}$  – age-0 fish in-stream survival (emergence to September 1<sup>st</sup>), year  $t - 2$ ,
- $\delta_{t-2}$  – age-0 fish winter survival (September 2<sup>nd</sup> – May 15<sup>th</sup>), year  $t - 2$ ,
- $\epsilon_{t-1}$  – age-1 fish survival (May 16<sup>th</sup> – May 15<sup>th</sup>), year  $t - 1$ , and
- $\theta_t$  – age-2 fish survival, year  $t$ .

It is assumed that a female that participates in the spawning run will deposit a clutch of eggs. The number of females in the spawning run is calculated as  $f_t \hat{N}_t$ , where  $f_t$  is the proportion of females captured during the spawning run in year  $t$ , and  $\hat{N}_t$  is the estimated spawning run population corrected for imperfect detection (e.g., Paterson 2013). Length-specific fecundity,  $\alpha_l$ , was estimated using data from Lund (1974) and Bishop (1971). Lund provided mean number of eggs and lengths by female length category; Bishop provided length and fecundity data from individuals. One of Bishop’s observations (13<sup>th</sup> observation) was excluded as an outlier. Total fecundity in year  $t$  is then  $F_t \alpha_l L_t$ , where  $L_t$  is mean female length in year  $t$ . Egg hatchability was taken from Lund’s (1974) work in Elk Lake. Hatchability varied from 0.04–0.12; the mean of these values ( $\bar{x} = 0.08$ ) was used for  $\beta$ .

Estimates of demographic rates were taken from published values for fish of similar life history, age, and size when empirical estimates were not otherwise available (Table 1). Maximum and mean survival rate values were obtained for model fitting. Mean Age-2 survival,  $\theta_t$ , was estimated using the upper confidence interval of annual survival for age-3 Red Rock Creek grayling (Paterson 2013). The upper confidence interval was selected because age-2 fish generally do not incur the risk of predation and physiological demands associated with spawning and, resultantly, likely have higher annual survival than age-3 fish. The maximum Age-2 survival rate was the highest annual adult survival rate estimated from available CMR data. Age-1 annual survival,  $\epsilon_{t-1}$ , and age-0 winter survival,  $\delta_{t-2}$ , were calculated by averaging published survival estimates for fish of similar life history, age, or size. Published survival estimates were transformed, when necessary, to account for differences between time intervals of published estimates and parameters of grayling models. Because no published

estimates applicable to age-0 in-stream survival,  $\gamma_{t-2}$ , were found, we calculated this rate for all years with adequate data using Equation 1 and the aforementioned age specific rates and solving for  $\gamma_{t-2}$ . Average age-0 in-stream survival was the average of the calculated rates among years and maximum age-0 in-stream survival was the highest annual value calculated.

**Table 1. Demographic estimates used for testing competing models of grayling response to winter habitat, spawning habitat, non-native predation, and spring hydrology.**

Species	Average survival rate (maximum survival rate)			
	$\gamma_{t-2}$	$\delta_{t-2}$	$\epsilon_{t-1}$	$\theta_t$
Arctic grayling <sup>1,2,3</sup>	<b>0.014 (0.035)</b> 15-100; 0-90 d 90 days	<b>0.25 (0.48)</b> 100-150; 90 d -1 y .75 year	<b>0.44 (0.68)</b> 153-211; 1-2 y 1 year	<b>0.74 (0.87)</b> 263-340; 2-3 y 1 year
Bull trout <sup>4</sup>	--	0.23 (0.38) 121-170; 2 y 1 year	--	--
Chinook salmon <sup>5</sup>	--	0.16 (0.48) 61-115; 90 d - 1.2 y 0.95 year	--	--
Bull trout <sup>6</sup>	--	0.09 (0.60) 121-170; 2y 1 year	0.45 (0.85) 171-220; 3y 1 year	--
Brown trout <sup>7</sup>	--	0.26 (0.47) 120-175; 0.5-1 y .75 year	0.43 (0.50) 200-305; 1-3 y 1 year	--

<sup>1</sup>Katzman 1998; <sup>2</sup>Mogen 1996; <sup>3</sup>Paterson 2014; <sup>4</sup>Bowerman, T. and P. Budy 2012; <sup>5</sup>Achord, S., R. Zabel, and B. Sanford 2007; <sup>6</sup>Al-Chokhachy, R. and P. Budy 2008; <sup>7</sup>Dieterman, D.J. and R.J.H. Hoxmeier 2011.

*Winter Habitat Model*—The influence of winter habitat on the grayling population would likely manifest itself as reduced survival of all-age grayling during years with widespread hypoxic conditions in Upper Red Rock Lake (e.g., Greenbank 1945). If the response of different age-class fish to winter habitat conditions is proportionally constant, e.g., poor winter conditions halve fish survival across all age classes, it is possible to estimate the relationship between all-age survival and winter conditions.

The influence of winter habitat conditions on grayling was quantified based on the minimum amount of winter habitat available between January and March, a period when hypoxic conditions can occur in Upper Red Rock Lake (Gangloff 1996, Davis 2016). Available winter habitat is defined as the area (ha) of water in Upper Red Rock Lake from January to March with  $\geq 4$  ppm dissolved oxygen and  $\geq 1$  m in depth (Davis 2016). Assuming species-specific density dependence, available winter habitat per fish,  $W_t$  (ha fish<sup>-1</sup>), is related to the area of suitable winter habitat,  $A_t$ , and the number of fish that entered the winter period,  $N_{w,t}$ .


$$W_t = \frac{A_t}{N_{w,t}} \quad (2)$$



The estimated number of spawning fish in Red Rock Creek in year  $t$  will be used as an index for  $N_{w,t}$ . Winter habitat will be related to the proportional change in all-age grayling survival using a saturating function (i.e., Holling type-II functional response) by

$$P_t = \frac{aW_t}{b+W_t}. \quad (3)$$

Parameters  $a$  and  $b$  determine how the realized proportion of maximum grayling survival is related to winter habitat conditions. Maximum realized proportion of grayling survival is  $a$ , and  $b$  represents the value of suitable winter habitat to an individual when the proportional change in survival is 50% of  $a$  (Hilborn and Mangel 1997). For example, if no reduction to survival occurs  $a = 1$ , i.e., grayling survive at their maximum age-class rates. To assess if the influence of available winter habitat is density independent,  $A_t$  will be substituted for  $W_t$  in Equation 3. Figure 3 shows a hypothetical situation where  $a = 1$ ,  $b = 10$ , and  $W_t$  varies from 0 to 0.50 ha fish<sup>-1</sup>.



**Figure 3.** Hypothetical relationship between the realized proportion of maximum grayling survival and the area of suitable winter habitat per fish in Upper Red Rock Lake based on a Holling type-II functional response.

The winter habitat model for grayling population dynamics and observation error, linking survival to winter habitat conditions, would then be:

$$N_{t+1} = N_t S_t P_t + F_{t-2} \alpha_{t-2} Y_{t-2} (\delta_{t-2} P_{t-2}) (\epsilon_{t-1} P_{t-1}) (\theta_t P_t), \quad (4)$$

$$N_{obs,t} = N_t V_t. \quad (5)$$


The number of adult fish surviving from year  $t$  to year  $t + 1$  is the product of the number of adults in year  $t$ , maximum annual survival ( $S_t$ ), and the realized proportion of maximum survival conditional on winter habitat conditions ( $P_t$ ). The number of potential recruits in year  $t$  is the number of age-2 fish, which is the product of the number of females  $t - 2$ , length-specific fecundity  $t - 2$ , the probability of an egg laid in year  $t - 2$  surviving until its first winter,  $Y_{t-2}$ , (the combined probabilities of egg ( $\beta$ ) and age-0 stream ( $\gamma$ ) survival), and maximum survival of age-0 winter ( $\delta$ ), and age-1 ( $\epsilon$ ) survival for cohort  $i$  multiplied by the estimated proportional influence of winter habitat on survival for each respective winter. The number of recruits in year  $t + 1$  is the product of the cohort in time  $t$ , second year survival ( $\theta_t$ ), and  $P_t$ . Substituting in demographic rates assumed fixed and constant (described above), gives the following equation for the winter habitat model

$$N_{t+1} = N_t * 0.74P_t + F_{t-2}\alpha_{t-2} * 0.0112 * (0.48P_{t-2})(0.68P_{t-1})(0.87P_t). \quad (6)$$

There are two components to the likelihood for this model, adult grayling annual abundance and survival. For the latter, apparent survival ( $\phi$ ) estimates for 1993–1996 (0.41, 95% CI = 0.24–0.66) and 2010–2013 (0.63, 95% CI = 0.53–0.74) are available (Paterson 2013). Estimates of  $\phi$  will be obtained annually using marked individuals.

*Spawning Habitat Model*—The relative quality of spawning habitat was hypothesized to influence cohort strength by its influence on egg ( $\beta$ ) and age-0 fish in-stream ( $\gamma$ ) survival. Low per capita area of suitable spawning habitat would lead to low egg and age-0 fish in-stream survival due to increased intra-specific competition for available spawning habitats, resulting in increased use of low suitability or unsuitable spawning habitat with lower intrinsic rates of egg and age-0 fish in-stream survival. Although degradation of spawning habitat is caused by the same mechanism (sedimentation) that degrades habitat for older fish, survival rates are most likely to be directly influenced in ages that are unable to avoid degraded habitat (i.e., eggs and fry).

The definition of suitable spawning habitat follows Hubert et al.’s (1985) functional relationships between suitability and percent fines and gravels in spawning riffles, where  $\leq 10\%$  fines is considered suitable, 11–50% fines represent linearly declining suitability, and  $> 50\%$  is unsuitable. Conversely,  $\geq 20\%$  gravel and rubble is considered suitable with  $< 20\%$  representing a linearly declining suitability (Figure 4). Thus, suitable spawning habitat can be characterized by having  $\leq 10\%$  fines and  $\geq 20\%$  gravel and rubble.



**Figure 4.** Predicted relationship between suitability of riverine Arctic grayling spawning habitat and a) percent fines (< 3 mm) in spawning areas and downstream riffles, and b) percent gravel and rubble (1.0–20.0 cm) in spawning areas (from Hubert et al. 1985).

The suitability threshold provided by Hubert et al. (1985) predicts the proposed asymptotic relationship between spawning area and recruitment. For example, at low population and high area of suitable spawning habitat, individuals would presumably all utilize the most suitable areas, resulting in maximum egg and age-0 fish in-stream survival, and number of recruits per individual. Further increases in suitable spawning habitat would not result in greater per capita recruitment. However, if population increased, and suitable spawning area per individual decreased, more individuals would spawn in less suitable habitats and an overall decrease in per capita recruitment would result as egg and age-0 fish in-stream survival declined.

Percent fines (particles < 3 mm) and gravel and rubble (1.0–20.0 cm) in riffles were estimated biennially using pebble count surveys. Each stream of interest was divided into reaches based on gross geomorphological characteristics and 1–5 representative sites were selected for sampling within each reach (Warren and Jaeger 2017). At each sampling site, four separate consecutive riffles were sampled following MT DEQ TMDL Sediment Assessment Methods (Kusnierz and Welch 2011). Cumulative percent fines are calculated for each sampled riffle.

Total area of suitable spawning habitat,  $A_t$ , was calculated and modeled considering 1) only habitat that has a suitability of 1.0 (i.e.,  $\leq 10\%$  fines and  $\geq 20\%$  gravel and rubble) and 2) weighted suitability of habitat based on observed percent fines and gravel and rubble following Hubert et al. (1985; Figure 4).

Habitat area per stream with suitability of 1.0,  $A_{t_s}$ , is riffle area per site with suitability = 1.0, divided by total site length, multiplied by reach length, summed across reaches within a stream (Equation 7).

$$\sum_{j=1}^n \frac{\sum_{i=1}^n \frac{\text{suitablerifflearea}(m^2)}{\text{sitelength}(m)}}{\text{numberofsites}} \times \text{reachlength}_j \quad (7)$$

Habitat area per stream with weighted suitability,  $A_{tw}$ , differs from  $A_{ts}$  in using the product of riffle suitability scores for percent fines and gravel and rubble estimated from the hypothesized suitability relationships of Hubert et al. (1985) instead of classifying riffle habitat as suitable (i.e., suitability = 1) or not (suitability = 0).


Area of suitable habitat ( $A_t$ ) was adjusted to account for effects of beaver dams and fragmentation. Habitat backwatered by beaver dams becomes unsuitable for spawning for at least the life of the beaver dam and the number and location of beaver dams varies among years. Each stream was annually surveyed and the total length of beaver dam backwaters was subtracted from each reach length when calculating  $A_t$ . The effects of fragmentation can range from incrementally reducing the likelihood of passage past a given location depending on daily conditions to completely precluding passage for that year. If passage is completely prevented then the area of upstream spawning habitat is functionally zero. If the probability of upstream passage is reduced then the area of available habitat is similarly reduced. To correct for the effects of fragmentation the area of suitable spawning habitats upstream of a barrier that prevents passage (i.e., probability of upstream passage is 0.0) was not included in calculation of  $A_t$ . Probability of passage at beaver dams was estimated based on assessment of relevant beaver dam characteristics within each reach (Cutting et al. 2019). Calculation of  $A_t$  was adjusted by multiplying the area of suitable habitat upstream of a beaver dam by the probability of passage at that dam. The effects of reduced passage probability were cumulatively considered. For example, the calculated value of  $A_t$  upstream of three beaver dams would be multiplied by the probability of a fish passing all three dams. Calculation of beaver dam effects was not required in 2017 and 2018 because of dam notching.

Assuming species-specific density dependence, availability of suitable spawning habitat per fish,  $H_t$  ( $m^2 \text{ fish}^{-1}$ ), is related to the area of suitable spawning habitat,  $A_t$ , and the number of spawning females,  $F_t$

$$H_t = \frac{A_t}{F_t}. \quad (8)$$

Spawning habitat was related to the product of egg and age-0 fish in-stream survival,  $R_t$ , using a saturating function (i.e., Holling type-II functional response) (Figure 5) by

$$R_t = \frac{aH_t}{b+H_t}. \quad (9)$$



**Figure 5.** Hypothetical relationship between grayling egg and age-0 fish in-stream survival,  $R$ , and area of suitable spawning habitat per female based on a Holling type-II functional response.

Parameters  $a$  and  $b$  determine how survival of eggs and age-0 fish are related to spawning habitat conditions. Maximum survival is  $a$ , and  $b$  represents the value of suitable spawning habitat when survival is 50% of  $a$  (Hilborn and Mangel 1997).

The spawning habitat model for grayling population dynamics and observation error, linking recruitment to spawning habitat conditions, is:


$$N_{t+1} = N_t S_t + F_{t-2} \alpha_{t-2} R_{t-2} \delta_{t-2} \epsilon_{t-1} \theta_t \quad (10)$$

$$N_{obs,t} = N_t V_t. \quad (11)$$

Adult grayling survival and total abundance year  $t$ , number of females year  $t - 2$  are obtained from sampling. Length-specific fecundity  $t - 2$  was estimated using data from Lund (1974) and Bishop (1971). Age-0 winter ( $\delta$ ), age-1 annual ( $\epsilon$ ), and age-2 annual ( $\theta$ ) fish survival were taken from published estimates for similar-aged salmonids or estimated for this grayling population (Table 1) and assumed to be constant among years. The product of survival estimates resulted in a value of 0.082, i.e.,  $\approx 8\%$  of age-0 fish that reach Upper Lake are predicted to survive through their second winter. There is only a single component to the likelihood for this model, adult grayling annual abundance estimates.

*Non-native Fish Model*—Non-native Yellowstone cutthroat trout (trout) were hypothesized to reduce survival of a grayling cohort prior to age-2, i.e., reduced age-0 through age-1 survival, via predation. To use the same model structure as the other hypotheses outlined above we considered grayling mortality ( $1 - \text{survival}$ ) instead of survival. This allows grayling mortality

to increase rapidly with increasing trout abundance up to a threshold at which mortality approaches an asymptote (Figure 6).



**Figure 6.** Hypothetical relationship between grayling age-0 and age-1 mortality for a cohort,  $Z$ , and concurrent winter abundance of adult Yellowstone cutthroat trout,  $C_t$ , based on a Holling type-II functional response.

Mortality of cohort  $i$  from hatching to age-1,  $Z_i (1 - \gamma_{t-2}\delta_{t-2}\epsilon_{t-1})$ , was asymptotically related to mean abundance of adult trout during the cohort's first two years. For example, mortality up to age-2 of a grayling cohort that hatched year  $t$  would be related to the mean abundance of adult trout in years  $t + 1$  and  $t + 2$ ,  $C_t$  as

$$Z_t = \frac{aC_t}{b+C_t}. \quad (12)$$

This results in a balance equation, relating grayling mortality to trout abundance, with the following form:

$$N_{t+1} = N_t S_t + F_{t-2} \alpha_{t-2} \beta_{t-2} Z_i \theta_t. \quad (13)$$

Adult trout abundance was annually estimated during spawning in Red Rock Creek by adding the number of fish 1) harvested by anglers, 2) removed at the fish weir, and 3) remaining in the system. Adult trout were experimentally removed from Red Rock Creek by culling fish at the weir (2013–2016) and liberalized angler harvest (2013–2017) to generate an adequately broad range of trout abundances to test this hypothesis. Number of fish harvested by anglers was estimated from catch cards corrected for non-reporting (Warren and Jaeger 2017). Cutthroat trout captured at the weir were sexed, length ( $\pm 1$  mm) and weight ( $\pm 1$  g) recorded, uniquely

marked with a floy tag and released upstream of the trap (2013–2017). Initial marking occurred in April; trout, marked and unmarked, encountered during electrofishing in May were recorded and used to estimate detection probability ( $p = \frac{m}{n}$ , where  $n$  is the number of trout marked in April and  $m$  is the number of marked trout recaptured in May). The number of trout remaining within Red Rock Creek was estimated as  $\frac{T}{p}$ , where  $T$  is the number of trout captured in May. Beginning in 2018, cutthroat trout abundance was estimated via two-pass electrofishing.

The enumeration of  $C_t$  likely provides a minimum estimate of the number of adult trout a given grayling cohort hatched year  $t$  was subjected to in years  $t + 1$  and  $t + 2$ . Because trout have a lower likelihood of being detected below the weir due to asynchronous timing of their spawning run and electrofishing surveys the number of fish remaining in the system will be underestimated. Therefore, the aforementioned overall abundance estimates represent an index of trout abundance that is less than actual abundance. It is possible that some adult trout present in the Upper Lake system do not ascend Red Rock Creek for spawning or complete spawning and return to Upper Lake prior to attempts to quantify their abundance. However,  $C_t$  is likely proportional to the number of adult trout present each year.

The *Non-native Fish* model does not differentiate between competition and predation, but will quantify the response of grayling to trout population reduction. Evidence for niche overlap between grayling and trout, where the potential for competition exists, occurs when trout are  $< 450$  mm in total length (Cutting et al. 2016). The management action being undertaken, Yellowstone cutthroat trout removal during spawning, is primarily removing larger ( $> 450$  mm) fish, which not only precludes a direct test of competition but also does not allow estimation of trout of the size class that potentially compete with grayling. Lastly, evidence for bottom-up regulation, e.g., low condition factor for either species observed during spawning, is lacking.

## Simulations, Predictions, and Model Weighting

Learning in the context of this adaptive management project occurs through the comparison of model predictions with reality (i.e., predicted grayling population vs. actual estimates of grayling population). Hypothesized drivers of grayling population dynamics are each represented by a model that links the driver to a specific life stage. A complete time series of observations (i.e., grayling and trout abundance, habitat characteristics) is needed to fit models so estimates of effects can be obtained and, subsequently, models can be used to make predictions. Delayed maturation of grayling (i.e., recruiting at age 3) results in needing three consecutive years of data to predict the number of grayling in a spawning population. The first time-series of data necessary to make a prediction was available in 2016, which was also the first population estimate that could be used to fit models. However, it is preferable to fit the models using several population estimates. To address this issue in the short-term, we conducted simulations to estimate the influence of each hypothesized driver of grayling population.

We conducted simulations to estimate response of grayling to 1) winter habitat, 2) spawning


habitat, and 3) cutthroat trout abundance. For each of the three models we simulated 1000 grayling populations for 15 years, using mean grayling population 1994–2018 ( $\hat{N} = 862$ ) as the first three population abundances (i.e.,  $N_t, N_{t+1}$ , and  $N_{t+2}$ ; necessary due to assumption of knife-edge recruitment at age-3). For each time step a random value was drawn from a normal distribution ( $X \sim N(\mu, \sigma^2)$ ) defined by the existing mean ( $\mu$ ) and standard deviation ( $\sigma$ ) from each variable (Figure 7). The distribution of winter habitat ( $W_t$ ) was defined by 8 values, with  $\mu = 96$  ha, and  $\sigma = 115$ . We assumed that a minimum of 5 ha was always available in the lake at the mouths of streams and spring heads; this precluded a complete die-off during a winter otherwise predicted to have 0 ha available habitat. A single estimate of weighted suitable spawning habitat was available; we used half that value for  $\mu$ , and assumed  $\sigma = 2$ , i.e.,  $A_{tw} \sim N(\mu = 0, \sigma = 2)$ . Finally, cutthroat trout abundance was drawn from  $N(\mu = 1132, \sigma = 964)$ , constrained to  $> 500$  trout. The *Non-native Fish Model* links mean trout abundance the first two years of a cohort’s life ( $C_t$ ) to mortality during that period (see **Models of System Dynamics** above). Therefore, we used the mean of two randomly drawn trout abundances to estimate  $C_t$ .

The common balance equation structure of models with saturating functions linking hypothesized population drivers to grayling demographic rates results in two parameters per model that mathematically describe the non-linear relationship. First,  $a$  represents maximum survival for the *Winter Habitat* and *Spawning Habitat* models. For the former,  $a$  is multiplied by the survival rate of each age-class to allow the influence of winter habitat to vary from none (i.e., grayling survival is 100% of expected age-class survival rates) to a winterkill event where survival  $\approx 0$ . For the *Spawning Habitat* model, maximum survival of egg ( $\beta$ ) and age-0 in-stream grayling ( $\gamma$ ),  $R_t$ , was set at 0.0042, the product of  $\beta = 0.12$  (Lund 1974) and  $\gamma = 0.035$  (maximum estimated survival based on back-calculations using existing demographic data). The complement of survival, mortality, is considered in the *Non-native Fish Model*. We set maximum combined mortality for first winter ( $\delta$ ), age-1 ( $\epsilon$ ), and age-2 ( $\theta$ ) grayling as 0.999, which results in a minimum survival of 0.001 for juvenile grayling, excluding mortality during egg and age-0 phases.

The second common parameter,  $b$ , defines the value of a given variable when survival is half of  $a$  (Hilborn and Mangel 1997) – the larger the value of  $b$ , the more sensitive survival is to the variable of interest.

We conducted simulations to estimate  $b$  for each model. Our convergence criterion was achieving a long-term simulated grayling population mean  $\pm 10$  individuals of the actual grayling mean, 1994–2018 ( $\hat{N} = 862$ ). Simulated population means did not include the first three values in the time series as these were set at the current grayling mean, 1994–current (see above).






**Figure 7.** Probability distributions used in grayling simulations to draw random a) minimum amount of suitable winter habitat (ha), b) amount of suitable spawning habitat (ha), and c) abundance of non-native cutthroat trout. Vertical dashed lines represent constraints, i.e.,  $> 5$  ha winter habitat and  $> 500$  trout.

Model weights were calculated using Baye’s formula, which allows adding new information (i.e., an updated comparison of predicted and observed grayling abundances) to existing information (i.e., existing model weights based on prior comparisons of predicted and observed grayling abundances). The model weight of model  $i$  in year  $t + 1$  given the observed data (i.e., response),  $p_{i,t+1}$ , is calculated as the prior model weight ( $p_t(model_i)$ ) multiplied by the probability of the observed data in  $t + 1$  given model  $i$  ( $P(response_{t+1}|model_i)$ ), divided by the total probability of all the models given the observed data ( $\sum_{j=1}^n p_t(model_j)P(response_{t+1}|model_j)$ ),

$$p_{i,t+1} = (model_i|response_{t+1}) = \frac{p_t(model_i)P(response_{t+1}|model_i)}{\sum_{j=1}^n p_t(model_j)P(response_{t+1}|model_j)} \quad (14)$$

We used observed values of  $W_t$ ,  $A_{ws}$  (uncorrected for barriers and constant across time),  $C_t$ , and  $\hat{b}$  from simulations for each model of system dynamics to predict the 2018 grayling spawning population in Red Rock Creek. Equivalent model weights (i.e., an uninformative prior) were used as model priors. Because enough data are not available to fit models and obtain likelihoods for  $P(response_{t+1}|model_i)$ , we used the probability of observing 387 grayling in 2018 given each model prediction, assuming a normal distribution with  $\mu =$  model predicted values and  $\sigma = 643$  (Figure 8). Model likelihoods based on maximum-likelihood estimation will be used for  $P(response_{t+1}|model_i)$  when a sufficient time-series is available for fitting models.




**Figure 8.** Normal probability density plots for *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models with  $\mu = 67, 92,$  and  $840,$  respectively. Standard deviation for all plots =  $643.$  Vertical dashed lines show the observed grayling spawning population,  $\hat{N} = 387,$  in 2018.


## RESULTS

### Grayling Abundance and Survival

The 2018 Red Rock Creek spawning population was 387 (95% CI = 168–682), similar to last year’s estimate of 176 fish (95% CI = 159–213)(Figure 9). The spawning population comprised primarily age-3 fish, consistent with the long-term age distribution of the spawning run. However, age-5 and older fish were underrepresented in 2018 compared to the long-term mean, a potential carry-over effect of the 2015–2016 winter mortality event (Figure 10).




**Figure 9.** Arctic grayling and Yellowstone cutthroat trout abundance estimates and 95% confidence intervals (grayling only) from Red Rock Creek, 2013–2018.



**Figure 10.** Age distribution of the spawning population of Arctic grayling in Red Rock Creek for a) all years of data (1950–2018, with missing years), and b) 2018.

## Winter Habitat

Suitable grayling winter habitat during 2018–2019 in Upper Red Rock Lake was surveyed 29 January, 2019. At that time, no suitable habitat was predicted in the lake (Fig. 11). This is only the second time no suitable habitat was predicted during 1995–2018 ( $n = 9$  winters, 21 sampling occasions). No suitable winter habitat was also estimated during the winter of 1995 (based on the water year starting October 1), the second year of a graduate project investigating grayling winter habitat use and distribution in Upper Lake (Figure 11; Gangloff 1996). Winter habitat suitability criteria, i.e., dissolved oxygen and water depth, were both highly variable during the period of record. For example, the area (ha) of Upper Lake with suitable dissolved oxygen and depth were 8–1016 ha, and 8–628, with mean values 272.5 (SD = 308.5) and 197.4 (SD = 196.4), respectively.



**Figure 11.** a) Extent of minimum area of suitable Arctic grayling winter habitat in Upper Red Rock Lake, 2018, b) 2019, c) annual estimate of minimum area of suitable habitat for water years 1995–2019, and d) grayling spawning population as a function of minimum area of suitable winter habitat for years when both were estimated.

## Spawning Habitat

Suitable spawning habitat was most recently quantified in 2017, with an estimated total area of suitable spawning habitat ( $A_{ts}$ ) of 0 ha, and weighted area of suitable habitat,  $A_{tw}$ , of 4 ha in Red Rock Creek (Table 2). All suitable spawning habitat was made available to grayling via beaver dam notching as part of the current AMP experiment. The increased per capita availability of suitable spawning habitat is hypothesized to increase egg ( $\beta$ ) and age-0 fish in-stream ( $\gamma$ ) survival. Spawning habitat will next be surveyed in 2019. The first year a response to increased availability of spawning habitat could be quantified is 2020, when the 2017 grayling cohort recruits.

**Table 2.** Estimated area (ha) of weighted suitable Arctic grayling spawning habitat (Hubert et al. 1985) by year and reach for Red Rock and Elk Springs creeks.

Creek	Reach	2013	2014	2015	2016	2017	2018
Red Rock Creek	Antelope Creek to Elk Lake Road	1.31	1.31	0.83	0.09	1.12	1.12
Red Rock Creek	Corral Creek to Antelope Creek	0.31	0.31	0.00	0.00	0.56	0.56
Red Rock Creek	Downstream of Battle Creek	0.04	0.04	0.03	0.03	0.00	0.00
Red Rock Creek	Elk Lake Road to Battle Creek	0.66	0.66	0.68	0.26	0.52	0.52
Red Rock Creek	Hellroaring Creek	0.00	0.00	0.00	0.00	0.42	0.42
Red Rock Creek	Huntsman	0.00	0.00	0.00	0.00	0.10	0.10
Red Rock Creek	Huntsman to Corral Creek	0.06	0.06	0.00	0.00	0.95	0.95
Elk Creek	Forks	0.00	0.00	0.00	0.00	0.11	0.11
Elk Creek	MacDonald Pond	0.00	0.00	0.00	0.00	0.00	0.00
Elk Creek	Restoration	0.00	0.00	0.00	0.00	0.20	0.20
	Total	2.38	2.38	1.54	0.38	3.97	3.97

## Simulations, Predictions, and Model Weighting

Simulations resulted in  $\hat{b}$  values of 0.0185, 0.217, and 1.225 for the *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models, respectively. These values of  $b$  resulted in simulated populations similar to observed dynamics 1994–2018, as measured by population mean during that period. Based on simulation results, the *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models predicted 67, 92, and 840 grayling, respectively, in the 2018 Red Rock Creek spawning population (Table 2). Updated model weights based on model predictions were 0.587, 0.311, and 0.103 for *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models, respectively. The *Spawning Habitat* model predicted the 2018 spawning run more precisely than the other two models (Figure 8; Table 1). Predicted grayling spawning population for 2019 are provided in Table 3.

**Table 3.** Arctic grayling spawning abundance model predictions, observed abundance, and relative model weights for 2018, and model predictions for 2019.

Model	2018 Prediction	Observed	Model Weights	2019 Prediction
Winter Habitat	67	387	0.587	21
Spawning Habitat	92	387	0.311	201
Non-native Fish	840	387	0.103	358

## Lituratione Cited

- Achord, S., R. Zabel, and B. Sanford. 2007. Migration timing, growth, and estimated parr-to-smolt survival rates of wild Snake River spring-summer chinook salmon from the Salmon River Basin, Idaho, to the lower Snake River. *Transactions of the American Fisheries Society* 136:142-154.
- Al-Chokhachy, R. and P. Budy. 2008. Demographic characteristics, population structure, and vital rates of a fluvial population of bull trout in Oregon. *Transactions of the American Fisheries Society* 137:1709-1722.
- Bishop, F. G. 1971. Observations on spawning habits and fecundity of the Arctic grayling. *Progressive Fish-Culturist* 33:12-19.
- Boltz, G. 2000. Fisheries investigations on the Red Rock Lakes National Wildlife Refuge, Montana. U.S. Fish and Wildlife Service, Bozeman, MT.
- Bowerman, T. and P. Budy. 2012. Incorporating movement patterns to improve survival estimates for bull trout. *North American Journal of Fisheries Management* 32: 1123-1136.
- Centennial Valley Historical Society, 2006. Centennial Valley: a journey through time 1820-1930. Vol. 1. Artcraft Printers, Butte, MT.
- Cutting, K. A., W. F. Cross, M. L. Anderson, and E. G. Reese. 2016. Seasonal change in trophic niche of adfluvial Arctic grayling (*Thymallus arcticus*) and coexisting fishes in a high-elevation lake system. *PLoS ONE* 11(5):e0156187.
- Cutting, K. A., J. M. Ferguson, M. L. Anderson, K. Cook, S. C. Davis, and R. Levine. 2018. Linking beaver dam affected flow dynamics to upstream passage of Arctic grayling. *Ecology and Evolution* 8:12905-12917.
- Deeds, J. F. and W. N. White. 1926. Water power and irrigation in the Jefferson River basin, Montana. U.S. Geologic Survey water-supply paper No. 580:41-116.
- Dieterman, D. J. and R. J. H. Hoxmeier. 2011. Demography of juvenile and adult brown trout in streams of southeastern Minnesota. *Transactions of the American Fisheries Society* 140:1642-1656.
- Gangloff, M. M. 1996. Winter habitat and distribution of Arctic grayling in Upper Red Rock Lake, Red Rock Lakes National Wildlife Refuge, Montana. Thesis, Montana State University, Bozeman, Montana.
- Greenbank, J. 1945. Limnological conditions in ice-covered lakes, especially as related to winter-kill of fish. *Ecological Monographs* 15:343-392.
- Hilborn, R. and M. Mangel. 1997. The ecological detective: confronting models with data. *Monographs in Population Biology* 28. Princeton University Press, Princeton, NJ.
- Hubert, W. A., R. S. Helzner, L. A. Lee, and P. C. Nelson. 1985. Habitat suitability index models and instream flow suitability curves: Arctic grayling riverine populations. Biological Report 82, Division of Biological Services, U.S. Fish and Wildlife Service, Washington D.C.

- Katzman, L. M. 1998. Effects of predation on status of Arctic grayling at Red Rock Lakes National Wildlife Refuge, Montana. Thesis, Montana State University, Bozeman, Montana.
- Kusnierz, P. and A. Welch. 2011. The Montana Department of Environmental Quality sediment assessment method: Considerations, physical and biological parameters, and decision making. 53p. Montana Dept. of Environmental Quality, Helena, MT.
- Laake, J. L. 2013. RMark: An R interface for analysis of capture-recapture data with MARK. AFSC Processed Report 2013-01, 25p. Alaska Fisheries Science Center, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Seattle WA.
- Lund, J. A. 1974. The reproduction of salmonids in the inlets of Elk Lake, Montana. Thesis, Montana State University, Bozeman, Montana.
- MAGWG (Montana Arctic grayling working group). In Press. Montana Arctic grayling conservation plan.
- MCA (Montana Code Annotated) 2000. United States Fish and Wildlife Service Red Rock Lakes-Montana Compact ratified. 85-20-801, Helena, MT.
- Mogen, J. T. 1996. Status and biology of the spawning population of Red Rock Lakes Arctic grayling. Thesis, Montana State University, Bozeman, Montana.
- MTFWP (Montana Fish, Wildlife & Parks). 1989. Application for reservations of water in the Missouri River basin above Fort Peck Dam, Volume 2, Reservation requests for waters above Canyon Ferry Dam. Unpublished Report, Montana Fish, Wildlife & Parks, Helena, MT.
- Mumma, S. A. 2010. A 20,000-yr-old record of vegetation and climate from Lower Red Rock Lake, Centennial Valley, southwestern Montana. Thesis. Montana State University, Bozeman, Montana.
- Nelson, P. H. 1954. Life history and management of the American Grayling (*Thymallus signifer tricolor*) in Montana. *Journal of Wildlife Management* 18:324-342.
- Oswald, R. A., S. Hochhalter, L. Rosenthal. 2008. Southwest Montana native fish research and conservation April 1, 2007–December 31, 2007. Montana Fish, Wildlife & Parks State Wildlife Grant Project Performance Report T-26-3, Helena, MT.
- Paterson, T. 2013. Estimation of the abundance and apparent survival of spawning Arctic grayling in Red Rock Creek; Red Rock Lakes National Wildlife Refuge. Thesis, Montana State University, Bozeman, Montana.
- Randall, L. C. 1978. Red Rock Lakes National Wildlife Refuge: an aquatic history 1899 - 1977. Unpublished report. U.S. Fish and Wildlife Service, Kalispell, Montana.
- Sonderegger, J. L. 1981. Geology and geothermal resources of the eastern Centennial Valley. Montana Geological Society, Field and Conference Symposium Guidebook to Southwest Montana.
- USFWS (U.S. Fish and Wildlife Service). 2009. Comprehensive Conservation Plan Red Rock Lakes National Wildlife Refuge. Lima, MT.



USFWS (U.S. Fish and Wildlife Service). 2014. Endangered and threatened wildlife and plants; revised 12-month finding on a petition to list the upper Missouri River distinct population segment of Arctic grayling as an endangered or threatened species; proposed rule. Federal Register, Vol. 79, No 161, 49384-49422.

Walters, C. 1986. Adaptive management of renewable resources. Blackburn Press, Caldwell, New Jersey.

Warren, J. M., and M. E. Jaeger. 2017. Centennial Valley Arctic grayling adaptive management plan. Available from Montana Fish, Wildlife and Parks, Dillon, MT.

White, G. C., and K. P. Burnham. 1999. Program MARK: Survival estimation from populations of marked animals. Bird Study 46 Supplement:120-138.