

Centennial Valley Arctic Grayling Adaptive Management Project Annual Report, 2019

25 March, 2020



EXECUTIVE SUMMARY

Arctic grayling (*Thymallus arcticus*) are a freshwater holarctic salmonid that were once widespread throughout the Upper Missouri River (UMR) drainage as a glacial relict population. One of the last endemic grayling populations remaining in the UMR drainage resides in the Centennial Valley (CV) of southwestern Montana. Spawning is largely limited to Red Rock, Corral, Elk Springs, and Odell creeks, with Red Rock Creek likely supporting 80-90% of annual spawning in the CV. It is presumed that most of the grayling population in the CV spends non-breeding portions of the year in Upper Red Rock Lake (Upper Lake). Red Rock Lakes National Wildlife Refuge (Refuge) encompasses Upper Lake, and nearly all of the currently occupied grayling spawning habitat within Red Rock, Elk Springs, and Odell creeks.

The estimated number of grayling in the 2019 Red Rock Creek spawning population was 154

(95% CI = 84–344), nearly unchanged from the previous year ($\hat{N} = 387$, 95% CI = 168–682; Figure 1). Suitable habitat the prior winter within Upper Lake (i.e., water depth below the ice ≥ 1 m and dissolved oxygen ≥ 4 ppm) reached a minimum during February sampling at an estimated 0 ha. This is the same area of suitable winter habitat experienced during the winter of 2015–2016 that led to the documented decline of spawning grayling from 1131 (95% CI = 1069–1210) in 2015 to 214 (95% CI = 161–321) in 2016. Area of winter habitat for the current year (2020) was ≈ 3 ha in January and February.

Suitable spawning habitat was quantified in 2019, with an estimated weighted area of suitable habitat (A_t) of 7 ha, in Red Rock and Elk Springs creeks (see **METHODS** below for description of variables). Barriers precluding grayling from accessing suitable spawning habitat, i.e., beaver dams, were notched prior to spawning again in 2019 (similar to 2017–2018) as part of the second hypothesis test of the Adaptive Management Plan.




Figure 1. Estimated annual Red Rock Creek Arctic grayling spawning population abundance and 95% confidence intervals (when available), 1994–2019. Confidence intervals are not available for some years prior to initiation of the Adaptive Management Plan.

The first management experiment conducted as part of the Adaptive Management Plan (AMP) was reducing non-native Yellowstone cutthroat trout (*Oncorhynchus clarkii bowieri*). Trout were captured and euthanized at a fish weir and fishing regulations were liberalized to 20 cutthroat trout per day (excluding a stream closure 15 May–14 June) on Red Rock Creek. Both actions were first implemented in 2013; the weir was operated through 2016 and liberalized harvest occurred through 2017. During 2013–2017 a total of 7149 cutthroat trout were removed from Red Rock Creek. Removal efforts peaked in 2014 at 2604 cutthroat

removed. The first year a grayling response to cutthroat removal could be quantified was 2016 when the 2013 grayling cohort recruited.

The second management experiment is exploring the relationship between spawning habitat and grayling recruitment. Efforts beginning in 2017 increased suitable habitat available for spawning by 1) ensuring fish passage at beaver dams ($n > 50$ in 2016) via notching, 2) restoration of an Elk Springs Creek channel that will provide direct connection to Upper Lake, and 3) restoration of spawning habitat at the head of Elk Springs Creek. This experiment will be undertaken for 4–5 years, similar to the cutthroat trout reduction. Increased per capita availability of spawning habitat is hypothesized to increase egg (β) and age-0 fish in-stream (γ) survival; the first year a possible response could be quantified is 2020, when the 2017 grayling cohort recruits.

Learning in the context of this project occurs through comparison of model predictions with reality (i.e., predicted grayling population vs. actual grayling population). Each hypothesized driver of grayling population dynamics, i.e., winter habitat, spawning habitat, and non-native fish, is represented by a model structured to estimate the driver's influence on a specific life stage. The *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models predicted 23, 245, and 358 grayling, respectively, in the 2019 Red Rock Creek spawning population. Model weights, i.e., relative support for a model given the data, were 0.556, 0.352, 0.092 for the *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models, respectively. The 2020 model-averaged prediction of spawning grayling abundance is 132 fish.

Authors:

Jeff Warren¹, Matt Jaeger², Andrew Gilham³, Kyle Cutting⁴, Lucas Bateman², Terrill Paterson², and Mike Duncan².

¹ U.S. Fish and Wildlife Service, Division of Science Resources,

² Montana Fish, Wildlife and Parks,

³ U.S. Fish and Wildlife Service, Montana Fish and Wildlife Conservation Office,

⁴ U.S. Fish and Wildlife Service, Red Rock Lakes National Wildlife Refuge.

INTRODUCTION

Montana Arctic grayling (*Thymallus arcticus*; hereafter grayling) were patchily distributed throughout the Upper Missouri River (UMR) drainage prior to the mid-1850s. This population declined to about 4% of their perceived historic distribution by the 1990s, which led to formal consideration for listing under the Endangered Species Act (USFWS 2014). One of the last populations of indigenous UMR grayling resides in the Centennial Valley (CV) of southwestern Montana. Grayling were historically distributed among at least a dozen CV streams and three lakes at presumably high abundances (Nelson 1954). Perceived distribution and abundance declined to historic lows sometime between the 1950s and mid-1990s, but have since improved, although large fluctuations in abundance still occur (USFWS 2014, MAGWG in press).

Currently, most of the CV grayling population spawns in Red Rock Creek and spends non-breeding portions of the year in Upper Red Rock Lake (Upper Lake) within Red Rock Lakes National Wildlife Refuge (Refuge; Figure 2). Over the past 70 years numerous hypotheses were posited regarding drivers of the CV grayling population, including 1) reduction and alteration of spawning habitat, 2) predation by, and competition with, non-native fishes, and 3) limited winter habitat. Although these hypotheses have been repeatedly proposed to explain population fluctuations, drivers of the population remain unclear.

Previous and ongoing research has focused on aspects of each hypothesis but has not linked them to demographic responses in grayling, which precludes inference regarding their role as population drivers. Resultantly, the most effective management and conservation approaches for CV grayling remain ambiguous, and selecting management actions can be contentious among and within agencies. This plan seeks to elucidate the relative effect of hypothesized drivers of CV grayling abundance to direct future management of this population. Determining the cause of previous population declines, *per se*, is not the primary issue of grayling conservation and management – finding an effective strategy to achieve population goals and prevent future declines is. In an effort to accomplish this, an adaptive management (AM) approach is being undertaken (Walters 1986). The Centennial Valley Arctic Grayling Adaptive Management Plan (AMP) embraces existing uncertainty regarding drivers of the grayling population in the CV to provide further understanding of important limiting factors and help guide management actions toward those that will have the most direct benefit to grayling.

Due to the initial level of structural uncertainty, and agency conflict regarding that uncertainty, the AMP was divided into two phases – a ‘management as experiment’ phase that emphasizes learning, i.e., reducing structural uncertainty (MacNab 1983, Walters 1986, Walters and Holling 1990), and an active adaptive management phase. The former was designed to explore grayling population response to hypothesized drivers that could be influenced via management actions. The latter will use the information gained in phase 1 to determine an optimal policy to inform annual management decisions (while still learning, but with less of an emphasis on learning).

STUDY AREA

The Centennial Valley of southwestern Montana is a high-elevation (ca. 2013 m) valley dominated by sagebrush steppe comprising *Artemisia* spp. shrub overstory and native bunchgrass understory (e.g., *Festuca* spp., *Nasella* spp., and *Hesperostipa* spp.). The valley is bounded on the north by the north-south trending Gravelly and Snowcrest mountain ranges and on the south by the east-west trending Centennial Mountains. Extensive wetlands exist throughout the CV, including a large shallow lake/wetland complex encompassed by Red Rock Lakes National Wildlife Refuge (Figure 2). The complex comprises Upper Red Rock, Lower Red Rock, and Swan lakes and associated palustrine emergent marsh dominated by seasonally-flooded sedge (*Carex* spp.). The complex is a remnant of Pleistocene Lake Centennial, a prehistoric lake that was believed to have formerly covered the valley floor to a depth of ca. 20 m (Mumma 2010). Upper Lake, the largest and deepest of the lakes, is ca. 1198 ha with a maximum depth of 2 m. The geologic (Sonderegger 1981; Centennial Valley Historical Society 2006), hydrologic (Deeds and White 1926, MTFWP 1989, MCA 2000), and fisheries (Nelson 1952, Randall 1978, Boltz 2000, Oswald et al. 2008) resources and contemporary administrative status (USFWS 2009) within the Centennial Valley are well described elsewhere.

The Centennial Valley includes all tributaries of the Red Rock River and their associated drainages upstream of Lima Dam (Figure 2). Most Upper Lake tributaries have their origins to the south along the eastern extent of the Centennial Mountains. Red Rock Creek, the largest of these tributaries, originates at an elevation of 2,562 m and flows north and west ca. 21 km to the northeast shore of Upper Lake. Elk Springs Creek originates from a series of springs south of Elk Lake and flows southwest, entering Upper Lake along the northeast shore. Red Rock River exits Upper Lake in the northwest corner, carrying water through the River Marsh and into the northwestern corner of Lower Lake. Red Rock River continues westward through the outlet of Lower Lake, ca.1.5 km west of where it enters the lake, leaving the CV near Lima, MT after passing through the 13 km long Lima Reservoir. Long Creek enters the Red Rock River 17 km downstream of the Lower Lake outlet and just upstream of Lima Reservoir.




Figure 2. Arctic grayling Adaptive Management Plan study area within the Centennial Valley of southwestern Montana.

METHODS

Abundance and Survival

Grayling and cutthroat trout were captured by mobile anode electrofishing (May and June) and a stationary weir (April–June) from 2013 to 2017 (see Paterson 2013 for further description). Beginning in 2018 only electrofishing surveys occurred. Grayling were uniquely marked with a visual implant (VI) tag, sexed, and length (± 1 mm) and weight (± 1 g) recorded. We used capture-mark-recapture (CMR) models implemented in program MARK (White and Burnham 1999) using the RMark package (Laake 2013) in R version 3.6.2 (R Development Core Team 2019) to estimate grayling abundance with closed population models and apparent survival (ϕ) with open population models concurrent with operation of the fish weir on Red Rock Creek. After cessation of the fish weir (2018), grayling and cutthroat estimates were obtained using electrofishing data and the Chapman method (Guy and Brown 2007). Apparent survival confounds permanent emigration and mortality; survival will be biased low if adult grayling fidelity to spawning streams is < 1 .

Models of System Dynamics

Spawning habitat, non-native fishes, and winter habitat have all been identified as potentially important drivers of grayling population dynamics in the upper CV. Each of these hypotheses is translated into a model, or set of models, to link hypothesized drivers and demographic

rates.

The annual abundance of spawning grayling is the product of demographic rates ranging from adult survival to the number of eggs deposited per female (fecundity) three years prior. All population models for spawning grayling share a common balance equation that allows prediction of annual abundance as a function of survival and recruitment processes:

$$N_{t+1} = N_t S_t + (F_{t-2} \alpha_{t-2} \beta_{t-2} \gamma_{t-2} \delta_{t-2} \epsilon_{t-1}) \theta_t \quad (1)$$

The number and survival of adult (i.e., reproductive age) grayling in year t is N_t and S_t , respectively. Assuming recruitment occurs with the age-3 cohort in year $t + 1$ (i.e., knife-edge recruitment at age-3), the number of potential age-2 recruits in year t is the product of:

- F_{t-2} – the number of females in the spawning run in year $t - 2$,
- α_{t-2} – length-specific fecundity rate, year $t - 2$,
- β_{t-2} – probability of an egg being fertilized and hatching, year $t - 2$,
- γ_{t-2} – age-0 fish in-stream survival (emergence to September 1st), year $t - 2$,
- δ_{t-2} – age-0 fish winter survival (September 2nd – May 15th), year $t - 2$,
- ϵ_{t-1} – age-1 fish survival (May 16th – May 15th), year $t - 1$, and
- θ_t – age-2 fish survival, year t .

It is assumed that a female that participates in the spawning run will deposit a clutch of eggs. The number of females in the spawning run is calculated as $f_t \hat{N}_t$, where f_t is the proportion of females captured during the spawning run in year t , and \hat{N}_t is the estimated spawning run population corrected for imperfect detection (e.g., Paterson 2013). Length-specific fecundity, α_l , was estimated using data from Lund (1974) and Bishop (1971). Lund provided mean number of eggs and lengths by female length category; Bishop provided length and fecundity data from individuals. One of Bishop’s observations (13th observation) was excluded as an outlier. Total fecundity in year t is then $F_t \alpha_l L_t$, where L_t is mean female length in year t . Egg hatchability was taken from Lund’s (1974) work in Elk Lake. Hatchability varied from 0.04–0.12; the mean of these values ($\bar{x} = 0.08$) was used for β .

Estimates of demographic rates were taken from published values for fish of similar life history, age, and size when empirical estimates were not otherwise available (Table 1). Maximum and mean survival rate values were obtained for model fitting. Mean Age-2 survival, θ_t , was estimated using the upper confidence interval of annual survival for age-3 Red Rock Creek grayling (Paterson 2013). The upper confidence interval was selected because age-2 fish generally do not incur the risk of predation and physiological demands associated with spawning and, resultantly, likely have higher annual survival than age-3 fish. The maximum Age-2 survival rate was the highest annual adult survival rate estimated from available CMR data. Age-1 annual survival, ϵ_{t-1} , and age-0 winter survival, δ_{t-2} , were calculated by averaging published survival estimates for fish of similar life history, age, or size. Published survival estimates were transformed, when necessary, to account for differences between time intervals of published estimates and parameters of grayling models. Because no published estimates applicable to age-0 in-stream survival, γ_{t-2} , were found, we calculated this rate

for all years with adequate data using Equation 1 and the aforementioned age specific rates and solving for γ_{t-2} . Average age-0 in-stream survival was the average of the calculated rates among years and maximum age-0 in-stream survival was the highest annual value calculated.

Table 1. Demographic estimates used for testing competing models of grayling response to winter habitat, spawning habitat, non-native predation, and spring hydrology.

| Species | Average survival rate (maximum survival rate) | | | |
|----------------------------------|---|--|--|--|
| | γ_{t-2} | δ_{t-2} | ϵ_{t-1} | θ_t |
| Arctic grayling ^{1,2,3} | 0.014 (0.035) 15-100; 0-90 d 90 days | 0.25 (0.48) 100-150; 90 d -1 y .75 year | 0.44 (0.68) 153-211; 1-2 y 1 year | 0.74 (0.87) 263-340; 2-3 y 1 year |
| Bull trout ⁴ | -- | 0.23 (0.38) 121-170; 2 y 1 year | -- | -- |
| Chinook salmon ⁵ | -- | 0.16 (0.48) 61-115; 90 d - 1.2 y 0.95 year | -- | -- |
| Bull trout ⁶ | -- | 0.09 (0.60) 121-170; 2y 1 year | 0.45 (0.85) 171-220; 3y 1 year | -- |
| Brown trout ⁷ | -- | 0.26 (0.47) 120-175; 0.5-1 y .75 year | 0.43 (0.50) 200-305; 1-3 y 1 year | -- |

¹Katzman 1998; ²Mogen 1996; ³Paterson 2014; ⁴Bowerman, T. and P. Budy 2012; ⁵Achord, S., R. Zabel, and B. Sanford 2007; ⁶Al-Chokhachy, R. and P. Budy 2008; ⁷Dieterman, D.J. and R.J.H. Hoxmeier 2011.

Winter Habitat Model—The influence of winter habitat on the grayling population would likely manifest itself as reduced survival of all-age grayling during years with widespread hypoxic conditions in Upper Red Rock Lake (e.g., Greenbank 1945). If the response of different age-class fish to winter habitat conditions is proportionally constant, e.g., poor winter conditions halve fish survival across all age classes, it is possible to estimate the relationship between all-age survival and winter conditions.

The influence of winter habitat conditions on grayling was quantified based on the minimum amount of winter habitat available between January and March, a period when hypoxic conditions can occur in Upper Red Rock Lake (Gangloff 1996, Davis 2016). Available winter habitat is defined as the area (ha) of water in Upper Red Rock Lake from January to March with ≥ 4 ppm dissolved oxygen and ≥ 1 m in depth (Davis 2016). Assuming species-specific density dependence, available winter habitat per fish, W_t (ha fish⁻¹), is related to the area of suitable winter habitat, A_t , and the number of fish that entered the winter period, $N_{w,t}$.

$$W_t = \frac{A_t}{N_{w,t}} \quad (2)$$

The estimated number of spawning fish in Red Rock Creek in year t will be used as an index for $N_{w,t}$. Winter habitat will be related to the proportional change in all-age grayling survival using a saturating function (i.e., Holling type-II functional response) by

$$P_t = \frac{aW_t}{b+W_t}. \quad (3)$$

Parameters a and b determine how the realized proportion of maximum grayling survival is related to winter habitat conditions. Maximum realized proportion of grayling survival is a , and b represents the value of suitable winter habitat to an individual when the proportional change in survival is 50% of a (Hilborn and Mangel 1997). For example, if no reduction to survival occurs $a = 1$, i.e., grayling survive at their maximum age-class rates. To assess if the influence of available winter habitat is density independent, A_t will be substituted for W_t in Equation 3. Figure 3 shows a hypothetical situation where $a = 1$, $b = 10$, and W_t varies from 0 to 0.50 ha fish⁻¹.




Figure 3. Hypothetical relationship between the realized proportion of maximum grayling survival and the area of suitable winter habitat per fish in Upper Red Rock Lake based on a Holling type-II functional response.

The winter habitat model for grayling population dynamics and observation error, linking survival to winter habitat conditions, would then be:

$$N_{t+1} = N_t S_t P_t + F_{t-2} \alpha_{t-2} Y_{t-2} (\delta_{t-2} P_{t-2}) (\epsilon_{t-1} P_{t-1}) (\theta_t P_t), \quad (4)$$

$$N_{obs,t} = N_t V_t. \quad (5)$$

The number of adult fish surviving from year t to year $t + 1$ is the product of the number of adults in year t , maximum annual survival (S_t), and the realized proportion of maximum survival conditional on winter habitat conditions (P_t). The number of potential recruits in year t is the number of age-2 fish, which is the product of the number of females $t - 2$, length-specific fecundity $t - 2$, the probability of an egg laid in year $t - 2$ surviving until its first winter, Y_{t-2} , (the combined probabilities of egg (β) and age-0 stream (γ) survival), and maximum survival of age-0 winter (δ), and age-1 (ϵ) survival for cohort i multiplied by the estimated proportional influence of winter habitat on survival for each respective winter. The number of recruits in year $t + 1$ is the product of the cohort in time t , second year survival (θ_t), and P_t . Substituting in demographic rates assumed fixed and constant (described above), gives the following equation for the winter habitat model

$$N_{t+1} = N_t * 0.74P_t + F_{t-2}\alpha_{t-2} * 0.0112 * (0.48P_{t-2})(0.68P_{t-1})(0.87P_t). \quad (6)$$

There are two components to the likelihood for this model, adult grayling annual abundance and survival. For the latter, apparent survival (ϕ) estimates for 1993–1996 (0.41, 95% CI = 0.24–0.66) and 2010–2013 (0.63, 95% CI = 0.53–0.74) are available (Paterson 2013). Estimates of ϕ will be obtained annually using marked individuals.

Spawning Habitat Model—The relative quality of spawning habitat was hypothesized to influence cohort strength by its influence on egg (β) and age-0 fish in-stream (γ) survival. Low per capita area of suitable spawning habitat would lead to low egg and age-0 fish in-stream survival due to increased intra-specific competition for available spawning habitats, resulting in increased use of low suitability or unsuitable spawning habitat with lower intrinsic rates of egg and age-0 fish in-stream survival. Although degradation of spawning habitat is caused by the same mechanism (sedimentation) that degrades habitat for older fish, survival rates are most likely to be directly influenced in ages that are unable to avoid degraded habitat (i.e., eggs and fry).

The definition of suitable spawning habitat follows Hubert et al.’s (1985) functional relationships between suitability and percent fines and gravels in spawning riffles, where $\leq 10\%$ fines is considered suitable, 11–50% fines represent linearly declining suitability, and $> 50\%$ is unsuitable. Conversely, $\geq 20\%$ gravel and rubble is considered suitable with $< 20\%$ representing a linearly declining suitability (Figure 4). Thus, the most suitable spawning habitat can be characterized by having $\leq 10\%$ fines and $\geq 20\%$ gravel and rubble.




Figure 4. Predicted relationship between suitability of riverine Arctic grayling spawning habitat and a) percent fines (< 3 mm) in spawning areas and downstream riffles, and b) percent gravel and rubble (1.0–20.0 cm) in spawning areas (from Hubert et al. 1985).

The suitability threshold provided by Hubert et al. (1985) predicts the proposed asymptotic relationship between spawning area and recruitment. For example, at low population and high area of suitable spawning habitat, individuals would presumably all utilize the most suitable areas, resulting in maximum egg and age-0 fish in-stream survival, and number of recruits per individual. Further increases in suitable spawning habitat would not result in greater per capita recruitment. However, if population increased, and suitable spawning area per individual decreased, more individuals would spawn in less suitable habitats and an overall decrease in per capita recruitment would result as egg and age-0 fish in-stream survival declined.

Percent fines (particles < 3 mm) and gravel and rubble (1.0–20.0 cm) in riffles were estimated biennially using pebble count surveys. Each stream of interest was divided into reaches based on gross geomorphological characteristics and 1–5 representative sites were selected for sampling within each reach (Warren and Jaeger 2017). At each sampling site, four separate consecutive riffles were sampled following MT DEQ TMDL Sediment Assessment Methods (Kusnierz and Welch 2011). Cumulative percent fines are calculated for each sampled riffle.

Total area of suitable spawning habitat, A_t , was calculated using weighted habitat suitability based on observed percent fines and gravel and rubble following Hubert et al. (1985; Figure 4). Empirical assessment of Arctic grayling egg survival at varying levels of sediment provides support for this approach (Anderson 2019). Riffle area (m^2 ; *riffle length (m) × riffle bankfull width (m)*) was multiplied by weighted riffle suitability to estimate area of suitable habitat by riffle. Area of suitable riffle habitat was then summed to obtain area of suitable spawning habitat by reach and stream.

To ensure all suitable spawning habitat was available to grayling as part of the current AMP treatment, beaver dams have been notched 2017–2019 to allow fish passage (i.e., probability

of passage = 1). During 2013–2016, area of suitable habitat (A_t) was adjusted to account for effects of beaver dams and fragmentation. Habitat backwatered by beaver dams becomes unsuitable for spawning for at least the life of the beaver dam and the number and location of beaver dams varies among years. Each stream was annually surveyed and the total length of beaver dam backwaters was subtracted from each reach length when calculating A_t . The effects of fragmentation can range from incrementally reducing the likelihood of passage past a given location depending on daily conditions to completely precluding passage for that year. If passage is completely prevented then the area of upstream spawning habitat is functionally zero. If the probability of upstream passage is reduced then the area of available habitat is similarly reduced. To correct for the effects of fragmentation the area of suitable spawning habitats upstream of a barrier that prevents passage (i.e., probability of upstream passage is 0.0) was not included in calculation of A_t . Probability of passage at beaver dams was estimated based on assessment of relevant beaver dam characteristics within each reach (Cutting et al. 2019). Calculation of A_t was adjusted by multiplying the area of suitable habitat upstream of a beaver dam by the probability of passage at that dam. The effects of reduced passage probability were cumulatively considered. For example, the calculated value of A_t upstream of three beaver dams would be multiplied by the probability of a fish passing all three dams.

Assuming species-specific density dependence, availability of suitable spawning habitat per fish, H_t ($\text{m}^2 \text{ fish}^{-1}$), is related to the area of suitable spawning habitat, A_t , and the number of spawning females, F_t

$$H_t = \frac{A_t}{F_t}. \quad (8)$$

Spawning habitat was related to the product of egg and age-0 fish in-stream survival, R_t , using a saturating function (i.e., Holling type-II functional response) (Figure 5) by

$$R_t = \frac{aH_t}{b+H_t}. \quad (9)$$




Figure 5. Hypothetical relationship between grayling egg and age-0 fish in-stream survival, R , and area of suitable spawning habitat per female based on a Holling type-II functional response.

Parameters a and b determine how survival of eggs and age-0 fish are related to spawning habitat conditions. Maximum survival is a , and b represents the value of suitable spawning habitat when survival is 50% of a (Hilborn and Mangel 1997).

The spawning habitat model for grayling population dynamics and observation error, linking recruitment to spawning habitat conditions, is:

$$N_{t+1} = N_t S_t + F_{t-2} \alpha_{t-2} R_{t-2} \delta_{t-2} \epsilon_{t-1} \theta_t \quad (10)$$

$$N_{obs,t} = N_t V_t. \quad (11)$$

Adult grayling survival and total abundance year t , number of females year $t - 2$ are obtained from sampling. Length-specific fecundity $t - 2$ was estimated using data from Lund (1974) and Bishop (1971). Age-0 winter (δ), age-1 annual (ϵ), and age-2 annual (θ) fish survival were taken from published estimates for similar-aged salmonids or estimated for this grayling population (Table 1) and assumed to be constant among years. The product of survival estimates resulted in a value of 0.082, i.e., $\approx 8\%$ of age-0 fish that reach Upper Lake are predicted to survive through their second winter. There is only a single component to the likelihood for this model, adult grayling annual abundance estimates.

Non-native Fish Model—Non-native Yellowstone cutthroat trout (trout) were hypothesized to reduce survival of a grayling cohort prior to age-2, i.e., reduced age-0 through age-1 survival, via predation. To use the same model structure as the other hypotheses outlined above we considered grayling mortality ($1 - \text{survival}$) instead of survival. This allows grayling mortality

to increase rapidly with increasing trout abundance up to a threshold at which mortality approaches an asymptote (Figure 6).




Figure 6. Hypothetical relationship between grayling age-0 and age-1 mortality for a cohort, Z , and concurrent winter abundance of adult Yellowstone cutthroat trout, C_t , based on a Holling type-II functional response.

Mortality of cohort i from hatching to age-1, $Z_i (1 - \gamma_{t-2}\delta_{t-2}\epsilon_{t-1})$, was asymptotically related to mean abundance of adult trout during the cohort's first two years. For example, mortality up to age-2 of a grayling cohort that hatched year t would be related to the mean abundance of adult trout in years $t + 1$ and $t + 2$, C_t as

$$Z_t = \frac{aC_t}{b+C_t}. \quad (12)$$

This results in a balance equation, relating grayling mortality to trout abundance, with the following form:

$$N_{t+1} = N_t S_t + F_{t-2} \alpha_{t-2} \beta_{t-2} Z_i \theta_t. \quad (13)$$

Adult trout abundance was annually estimated during spawning in Red Rock Creek by adding the number of fish 1) harvested by anglers, 2) removed at the fish weir, and 3) remaining in the system. Adult trout were experimentally removed from Red Rock Creek by culling fish at the weir (2013–2016) and liberalized angler harvest (2013–2017) to generate an adequately broad range of trout abundances to test this hypothesis. Number of fish harvested by anglers was estimated from catch cards corrected for non-reporting (Warren and Jaeger 2017). Cutthroat trout captured at the weir were sexed, length (± 1 mm) and weight (± 1 g) recorded, uniquely

marked with a floy tag and released upstream of the trap (2013–2017). Initial marking occurred in April; trout, marked and unmarked, encountered during electrofishing in May were recorded and used to estimate detection probability ($p = \frac{m}{n}$, where n is the number of trout marked in April and m is the number of marked trout recaptured in May). The number of trout remaining within Red Rock Creek was estimated as $\frac{T}{p}$, where T is the number of trout captured in May. Beginning in 2018, cutthroat trout abundance was estimated via two-pass electrofishing and the Chapman method.

The enumeration of C_t likely provides a minimum estimate of the number of adult trout a given grayling cohort hatched year t was subjected to in years $t + 1$ and $t + 2$. Because trout have a lower likelihood of being detected below Elk Lake Road due to asynchronous timing of their spawning run and electrofishing surveys, the number of fish remaining in the system will be underestimated. Therefore, the aforementioned overall abundance estimates represent an index of trout abundance that is less than actual abundance. It is possible that some adult trout present in the Upper Lake system do not ascend Red Rock Creek for spawning or complete spawning and return to Upper Lake prior to attempts to quantify their abundance. However, C_t is likely proportional to the number of adult trout present each year.

The *Non-native Fish* model does not differentiate between competition and predation, but will quantify the response of grayling to trout population reduction. Evidence for niche overlap between grayling and trout, where the potential for competition exists, occurs when trout are < 450 mm in total length (Cutting et al. 2016). The management action being undertaken, Yellowstone cutthroat trout removal during spawning, is primarily removing larger (> 450 mm) fish, which not only precludes a direct test of competition but also does not allow estimation of trout of the size class that potentially compete with grayling. Lastly, evidence for bottom-up regulation, e.g., low condition factor for either species observed during spawning, is lacking.

Simulations, Predictions, and Model Weighting

Learning in the context of this adaptive management project occurs through the comparison of model predictions with reality (i.e., predicted grayling population vs. actual estimates of grayling population). Hypothesized drivers of grayling population dynamics are each represented by a model that links the driver to a specific life stage. A complete time series of observations (i.e., grayling and trout abundance, habitat characteristics) is needed to fit models so estimates of effects can be obtained and, subsequently, models can be used to make predictions. Delayed maturation of grayling (i.e., recruiting at age 3) results in needing three consecutive years of data to predict the number of grayling in a spawning population. The first time-series of data necessary to make a prediction was available in 2016, which was also the first population estimate that could be used to fit models. However, it is preferable to fit the models using several population estimates. To address this issue in the short-term, we conducted simulations to estimate the influence of each hypothesized driver of grayling population.

We conducted simulations to estimate response of grayling to 1) winter habitat, 2) spawning habitat, and 3) cutthroat trout abundance. For each of the three models we simulated 1000

grayling populations for 15 years, using mean grayling population 1994–2019 ($\hat{N} = 829$) as the first three population abundances (i.e., N_t, N_{t+1} , and N_{t+2} ; necessary due to assumption of knife-edge recruitment at age-3). For each time step a random value was drawn from a normal distribution ($X \sim N(\mu, \sigma^2)$) defined by the existing mean (μ) and standard deviation (σ) from each variable (Figure 7). The distribution of winter habitat (W_t) was defined by 9 values, with $\mu = 114$ ha, and $\sigma = 148$. We assumed that a minimum of 5 ha was always available in the lake at the mouths of streams and spring heads; this precluded a complete die-off during a winter otherwise predicted to have 0 ha available habitat. Estimates of weighted suitable spawning habitat, 2013–2019, were used to estimate mean (μ) and variance (σ) spawning habitat values for simulations ($A_w \sim N(\mu = 3.26, \sigma = 2.17)$). Finally, cutthroat trout abundance was drawn from $N(\mu = 1088, \sigma = 947)$, constrained to > 500 trout. The *Non-native Fish Model* links mean trout abundance the first two years of a cohort’s life (C_t) to mortality during that period (see **Models of System Dynamics** above). Therefore, we used the mean of two randomly drawn trout abundances to estimate C_t .

The common balance equation structure of models with saturating functions linking hypothesized population drivers to grayling demographic rates results in two parameters per model that mathematically describe the non-linear relationship. First, a represents maximum survival for the *Winter Habitat* and *Spawning Habitat* models. For the former, a is multiplied by the survival rate of each age-class to allow the influence of winter habitat to vary from none (i.e., grayling survival is 100% of expected age-class survival rates) to a winterkill event where survival ≈ 0 . For the *Spawning Habitat* model, maximum survival of egg (β) and age-0 in-stream grayling (γ), R_t , was set at 0.0042, the product of $\beta = 0.12$ (Lund 1974) and $\gamma = 0.035$ (maximum estimated survival based on back-calculations using existing demographic data). The complement of survival, mortality, is considered in the *Non-native Fish Model*. We set maximum combined mortality for first winter (δ), age-1 (ϵ), and age-2 (θ) grayling as 0.999, which results in a minimum survival of 0.001 for juvenile grayling, excluding mortality during egg and age-0 phases.

The second common parameter, b , defines the value of a given variable when survival is half of a (Hilborn and Mangel 1997) – the larger the value of b , the more sensitive survival is to the variable of interest.

We conducted simulations to estimate b for each model. Our convergence criterion was achieving a long-term simulated grayling population mean ± 10 individuals of the actual grayling mean, 1994–2019 ($\hat{N} = 829$). Simulated population means did not include the first three values in the time series as these were set at the current grayling mean, 1994–current (see above).




Figure 7. Probability distributions used in grayling simulations to draw random a) minimum amount of suitable winter habitat (ha), b) amount of suitable spawning habitat (ha), and c) abundance of non-native cutthroat trout. Vertical dashed lines represent constraints, i.e., > 5 ha winter habitat and > 500 trout.

Model weights were calculated using Baye’s formula, which allows adding new information (i.e., an updated comparison of predicted and observed grayling abundances) to existing information (i.e., existing model weights based on prior comparisons of predicted and observed grayling abundances). The model weight of model i in year $t + 1$ given the observed data (i.e., response), $p_{i,t+1}$, is calculated as the prior model weight ($p_t(model_i)$) multiplied by the probability of the observed data in $t + 1$ given model i ($P(response_{t+1}|model_i)$), divided by the total probability of all the models given the observed data ($\sum_{j=1}^n p_t(model_j)P(response_{t+1}|model_j)$),

$$p_{i,t+1} = (model_i|response_{t+1}) = \frac{p_t(model_i)P(response_{t+1}|model_i)}{\sum_{j=1}^n p_t(model_j)P(response_{t+1}|model_j)} \quad (14)$$

We used observed values of W_t , A_t , C_t , and \hat{b} from simulations for each model of system dynamics to predict the 2019 grayling spawning population in Red Rock Creek. Prior-year’s (2018) model weights were used as model priors. Because enough data are not available to fit models and obtain likelihoods for $P(response_{t+1}|model_i)$, we used the probability of observing 154 grayling in 2019 given each model prediction, assuming a normal distribution with $\mu =$ model predicted values and $\sigma = 645$ (Figure 8). Model likelihoods based on maximum-likelihood estimation will be used for $P(response_{t+1}|model_i)$ when a sufficient time-series is available for fitting models.




Figure 8. Normal probability density plots for *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models with $\mu = 23, 245,$ and $358,$ respectively. Standard deviation for all plots = 645. Vertical dashed lines show the observed grayling spawning population, $\hat{N} = 154,$ in 2019.

RESULTS

Abundance and Survival

The 2019 Red Rock Creek grayling spawning population was 154 fish (95% CI = 84–344), similar to last year’s estimate of 387 grayling (95% CI = 168–682)(Figure 9). The spawning population comprised primarily age-5 fish, which is inconsistent with the long-term age distribution of the spawning run. Age-3 fish were underrepresented in 2019 compared to the long-term mean, a possible carry-over effect of four consecutive potential winter mortality events (Figure 10). The estimated spawning population of cutthroat trout was 440 fish (95% CI = 394–504), an approximate decline of 52% from 2018. This decline indicates cutthroat trout may be negatively affected by winter conditions in Upper Lake similar to grayling. Any evidence of this relationship during the first four years of the AMP would have been confounded with, and masked by, the trout removal taking place at that time. Moreover, the correlation coefficient between grayling and trout spawning populations 2012–2019 is $r(6) = 0.74$ ($p = 0.03$).




Figure 9. Arctic grayling and Yellowstone cutthroat trout abundance estimates and 95% confidence intervals (grayling only) from Red Rock Creek, 2013–2019.




Figure 10. Age distribution of the spawning population of Arctic grayling in Red Rock Creek for a) all years of data (1950–2019, with missing years), and b) 2019.

Winter Habitat

Suitable grayling winter habitat during 2019–2020 in Upper Red Rock Lake was surveyed 21 January and 28 February, 2020. During both sampling occasions ≈ 3 ha of suitable habitat was estimated in the lake (Fig. 11). Winter habitat suitability criteria, i.e., dissolved oxygen and water depth, were both highly variable during the period of record. For example, the area (ha) of Upper Lake with suitable dissolved oxygen and depth were 7–1023 ha, and 8–628, with mean values 324.8 (SD = 330.6) and 184.6 (SD = 193.2), respectively.




Figure 11. a) Extent of minimum area of suitable Arctic grayling winter habitat in Upper Red Rock Lake, 2019, b) 2020, c) annual estimate of minimum area of suitable habitat for water years 1995–2020, and d) grayling spawning population as a function of minimum area of suitable winter habitat for years when both were estimated.

Spawning Habitat

Suitable spawning habitat was most recently quantified in 2019, with an estimated weighted area of suitable habitat, A_t , of 7 ha in Red Rock Creek (Table 2, Appendix). Area of suitable spawning habitat increased 60% between 2017 and 2019. This increase was coincident with

above average flows on Red Rock Creek; days above 67% of bankfull discharge ($c67bf = 38$) were 179% of the long-term mean. Fines within riffles are mobilized by stream discharges greater than 67% of bankfull (Mueller et al. 2005), providing a mechanism for the observed improvement in grayling spawning habitat in 2019 (Hubbert et al. 1985). All suitable spawning habitat was made available to grayling via beaver dam notching as part of the current AMP experiment. The increased per capita availability of suitable spawning habitat is hypothesized to increase egg (β) and age-0 fish in-stream (γ) survival. The first year a response to increased availability of spawning habitat could be quantified is 2020, when the 2017 grayling cohort recruits. Spawning habitat will next be surveyed in 2021.

Table 2. Estimated area (ha) of weighted suitable Arctic grayling spawning habitat (Hubbert et al. 1985) by year and stream for Red Rock and Elk Springs creeks.

| Year | Red Rock Creek | Elk Springs Creek | Total |
|------|----------------|-------------------|-------|
| 2013 | 2.55 | 0.00 | 2.55 |
| 2014 | 2.55 | 0.00 | 2.55 |
| 2015 | 1.63 | 0.00 | 1.63 |
| 2016 | 0.38 | 0.00 | 0.38 |
| 2017 | 4.04 | 0.33 | 4.37 |
| 2018 | 4.04 | 0.33 | 4.37 |
| 2019 | 6.88 | 0.10 | 6.98 |

Simulations, Predictions, and Model Weighting

Simulations resulted in \hat{b} values of 0.0185, 0.011, and 1.225 for the *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models, respectively. These values of b resulted in simulated populations similar to observed dynamics 1994–2019, as measured by population mean during that period. Based on simulation results, the *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models predicted 23, 245, and 358 grayling, respectively, in the 2019 Red Rock Creek spawning population (Table 3). Updated model weights based on model predictions were 0.556, 0.352, and 0.092 for *Winter Habitat*, *Spawning Habitat*, and *Non-native Fish* models, respectively. The *Spawning Habitat* model predicted the 2019 spawning run more precisely than the other two models (Figure 8; Table 1). Each model’s predicted grayling spawning population for 2020 are provided in Table 3. The 2020 model-averaged prediction of spawning grayling abundance is 132 fish.

Table 3. Arctic grayling spawning abundance model predictions, observed abundance, and relative model weights for 2019, and model predictions for 2020.

| Model | 2019 Prediction | Observed | Model Weights | 2020 Prediction |
|------------------|-----------------|----------|---------------|-----------------|
| Winter Habitat | 23 | 154 | 0.556 | 26 |
| Spawning Habitat | 245 | 154 | 0.352 | 260 |
| Non-native Fish | 358 | 154 | 0.092 | 206 |

Lituratione Cited

- Achord, S., R. Zabel, and B. Sanford. 2007. Migration timing, growth, and estimated parr-to-smolt survival rates of wild Snake River spring-summer chinook salmon from the Salmon River Basin, Idaho, to the lower Snake River. *Transactions of the American Fisheries Society* 136:142-154.
- Al-Chokhachy, R. and P. Budy. 2008. Demographic characteristics, population structure, and vital rates of a fluvial population of bull trout in Oregon. *Transactions of the American Fisheries Society* 137:1709-1722.
- Anderson, M. 2019. Sediment effects on Arctic grayling (*Thymallus arcticus*) early life history in Montana – a field experiment and literature review. Project completion report to Montana Fish, Wildlife, and Parks, Helena, MT.
- Bishop, F. G. 1971. Observations on spawning habits and fecundity of the Arctic grayling. *Progressive Fish-Culturist* 33:12-19.
- Boltz, G. 2000. Fisheries investigations on the Red Rock Lakes National Wildlife Refuge, Montana. U.S. Fish and Wildlife Service, Bozeman, MT.
- Bowerman, T. and P. Budy. 2012. Incorporating movement patterns to improve survival estimates for bull trout. *North American Journal of Fisheries Management* 32: 1123-1136.
- Centennial Valley Historical Society, 2006. Centennial Valley: a journey through time 1820-1930. Vol. 1. Artcraft Printers, Butte, MT.
- Cutting, K. A., W. F. Cross, M. L. Anderson, and E. G. Reese. 2016. Seasonal change in trophic niche of adfluvial Arctic grayling (*Thymallus arcticus*) and coexisting fishes in a high-elevation lake system. *PLoS ONE* 11(5):e0156187.
- Cutting, K. A., J. M. Ferguson, M. L. Anderson, K. Cook, S. C. Davis, and R. Levine. 2018. Linking beaver dam affected flow dynamics to upstream passage of Arctic grayling. *Ecology and Evolution* 8:12905-12917.
- Deeds, J. F. and W. N. White. 1926. Water power and irrigation in the Jefferson River basin, Montana. U.S. Geologic Survey water-supply paper No. 580:41-116.
- Dieterman, D. J. and R. J. H. Hoxmeier. 2011. Demography of juvenile and adult brown trout in streams of southeastern Minnesota. *Transactions of the American Fisheries Society* 140:1642-1656.
- Gangloff, M. M. 1996. Winter habitat and distribution of Arctic grayling in Upper Red Rock Lake, Red Rock Lakes National Wildlife Refuge, Montana. Thesis, Montana State University, Bozeman, Montana.
- Greenbank, J. 1945. Limnological conditions in ice-covered lakes, especially as related to winter-kill of fish. *Ecological Monographs* 15:343-392.
- Guy, C. S. and M. L. Brown. 2007. Analysis and interpretation of freshwater fisheries data. American Fisheries Society, Bethesda, MD.

- Hilborn, R. and M. Mangel. 1997. The ecological detective: confronting models with data. Monographs in Population Biology 28. Princeton University Press, Princeton, NJ.
- Hubert, W. A., R. S. Helzner, L. A. Lee, and P. C. Nelson. 1985. Habitat suitability index models and instream flow suitability curves: Arctic grayling riverine populations. Biological Report 82, Division of Biological Services, U.S. Fish and Wildlife Service, Washington D.C.
- Katzman, L. M. 1998. Effects of predation on status of Arctic grayling at Red Rock Lakes National Wildlife Refuge, Montana. Thesis, Montana State University, Bozeman, Montana.
- Kusnierz, P. and A. Welch. 2011. The Montana Department of Environmental Quality sediment assessment method: Considerations, physical and biological parameters, and decision making. 53p. Montana Dept. of Environmental Quality, Helena, MT.
- Laake, J. L. 2013. RMark: An R interface for analysis of capture-recapture data with MARK. AFSC Processed Report 2013-01, 25p. Alaska Fisheries Science Center, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Seattle WA.
- Lund, J. A. 1974. The reproduction of salmonids in the inlets of Elk Lake, Montana. Thesis, Montana State University, Bozeman, Montana.
- MAGWG (Montana Arctic grayling working group). In Press. Montana Arctic grayling conservation plan.
- MCA (Montana Code Annotated) 2000. United States Fish and Wildlife Service Red Rock Lakes-Montana Compact ratified. 85-20-801, Helena, MT.
- Mogen, J. T. 1996. Status and biology of the spawning population of Red Rock Lakes Arctic grayling. Thesis, Montana State University, Bozeman, Montana.
- MTFWP (Montana Fish, Wildlife & Parks). 1989. Application for reservations of water in the Missouri River basin above Fort Peck Dam, Volume 2, Reservation requests for waters above Canyon Ferry Dam. Unpublished Report, Montana Fish, Wildlife & Parks, Helena, MT.
- Mumma, S. A. 2010. A 20,000-yr-old record of vegetation and climate from Lower Red Rock Lake, Centennial Valley, southwestern Montana. Thesis. Montana State University, Bozeman, Montana.
- Nelson, P. H. 1954. Life history and management of the American Grayling (*Thymallus signifer tricolor*) in Montana. *Journal of Wildlife Management* 18:324-342.
- Oswald, R. A., S. Hochhalter, L. Rosenthal. 2008. Southwest Montana native fish research and conservation April 1, 2007–December 31, 2007. Montana Fish, Wildlife & Parks State Wildlife Grant Project Performance Report T-26-3, Helena, MT.
- Paterson, T. 2013. Estimation of the abundance and apparent survival of spawning Arctic grayling in Red Rock Creek; Red Rock Lakes National Wildlife Refuge. Thesis, Montana State University, Bozeman, Montana.
- Randall, L. C. 1978. Red Rock Lakes National Wildlife Refuge: an aquatic history 1899 - 1977. Unpublished report. U.S. Fish and Wildlife Service, Kalispell, Montana.

Sonderegger, J. L. 1981. Geology and geothermal resources of the eastern Centennial Valley. Montana Geological Society, Field and Conference Symposium Guidebook to Southwest Montana.

USFWS (U.S. Fish and Wildlife Service). 2009. Comprehensive Conservation Plan Red Rock Lakes National Wildlife Refuge. Lima, MT.

USFWS (U.S. Fish and Wildlife Service). 2014. Endangered and threatened wildlife and plants; revised 12-month finding on a petition to list the upper Missouri River distinct population segment of Arctic grayling as an endangered or threatened species; proposed rule. Federal Register, Vol. 79, No 161, 49384-49422.

Walters, C. 1986. Adaptive management of renewable resources. Blackburn Press, Caldwell, New Jersey.

Warren, J. M., and M. E. Jaeger. 2017. Centennial Valley Arctic grayling adaptive management plan. Available from Montana Fish, Wildlife and Parks, Dillon, MT.

White, G. C., and K. P. Burnham. 1999. Program MARK: Survival estimation from populations of marked animals. Bird Study 46 Supplement:120-138.

Appendix. Estimated area (ha) of weighted suitable Arctic grayling spawning habitat (Hubert et al. 1985) in Red Rock and Elk Spring creeks by year and reach.

| Creek | Reach | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 |
|----------------|---------------------------------|------|------|------|------|------|------|------|
| Red Rock Creek | Antelope Creek to Elk Lake Road | 1.47 | 1.47 | 0.93 | 0.11 | 1.26 | 1.26 | 3.03 |
| Red Rock Creek | Corral Creek to Antelope Creek | 0.34 | 0.34 | 0.00 | 0.00 | 0.59 | 0.59 | 0.48 |
| Red Rock Creek | Downstream of Battle Creek | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| Red Rock Creek | Elk Lake Road to Battle Creek | 0.68 | 0.68 | 0.70 | 0.27 | 0.53 | 0.53 | 1.19 |
| Red Rock Creek | Hellroaring Creek | 0.00 | 0.00 | 0.00 | 0.00 | 0.44 | 0.44 | 0.47 |
| Red Rock Creek | Huntsman | 0.00 | 0.00 | 0.00 | 0.00 | 0.10 | 0.10 | 0.28 |
| Red Rock Creek | Huntsman to Corral Creek | 0.06 | 0.06 | 0.00 | 0.00 | 1.12 | 1.12 | 1.43 |
| Elk Creek | Forks | 0.00 | 0.00 | 0.00 | 0.00 | 0.11 | 0.11 | 0.03 |
| Elk Creek | Restoration | 0.00 | 0.00 | 0.00 | 0.00 | 0.22 | 0.22 | 0.07 |
| | Total | 2.55 | 2.55 | 1.63 | 0.38 | 4.37 | 4.37 | 6.98 |