

Vital rates, limiting factors and monitoring methods for moose in Montana



Federal Aid in Wildlife Restoration Grant W-157-R-7
Annual report, September 1, 2019

Nick DeCesare

Research Wildlife Biologist, MFWP
3201 Spurgin Rd. | Missoula, MT 59804
406-542-5558 | ndecesare@mt.gov

Jesse Newby

Research Technician, MFWP
490 N. Meridian | Kalispell, MT 59901
406-751-4588 | jnewby@mt.gov



State: Montana

Agency: Fish, Wildlife & Parks

Grant: Montana Shiras Moose Study

Grant number: W-157-R-7

Time period: 1 July, 2018 – 30 June, 2019

FUNDING PARTNERS:

Safari Club International Foundation:
Conservation Grants, 2016, 2018



Note: All results should be considered preliminary and subject to change; please contact the authors before citing or referencing these data.

Background and summary

Concern has arisen in recent years over widespread declines of North American moose (*Alces alces*) populations along the southern extent of their range. Populations in Montana appear to have declined since the 1990's, as evidenced by aerial survey trends and hunter harvest statistics. While declining populations have clear implications for hunting opportunity, moose management in Montana also suffers from a lack of rigorous data and methods with which to monitor population trends and prescribe actions.

In 2013, Montana Fish, Wildlife, & Parks (MFWP) began a 10-year study designed to improve our understanding of: 1) cost-effective means to monitor statewide moose populations, and 2) the current status and trends of moose populations and the relative importance of factors influencing moose vital rates and limiting population growth (including predators, disease, habitat, and weather). We are using a mechanistic approach to hierarchically assess which factors are drivers of moose vital rates (e.g., adult survival, pregnancy, calf survival), and ultimately influence annual growth of moose populations.

This document is the 7th annual report produced as part of this work. This report contains preliminary results from a subset of our work, including results from the first 6 biological years of moose research and monitoring. All results should be considered preliminary as both data collection and analyses are works in progress.

We continue to explore whether monitoring moose with hunter observations may offer a promising new approach to gathering statewide data. After collecting >4,300 statewide moose sighting locations per year during 2012–2016 through annual hunter phone surveys, we have moved analyses of these data from an occupancy-based (presence-absence) framework into an abundance-based (counts) framework. Results from this work will be finalized during FY20.

Moose vital rates measured with radio-collar studies currently indicate stable to increasing average annual population trends in all 3 study areas (Cabinet-Salish, Big Hole Valley and Rocky Mountain Front). These estimated trends are largely driven by adult female survival rates, which have been variable year by year, but generally have averaged moderate rates of 0.86–0.89 over the full study period. Calf survival appears to have less influence on the overall trajectory of the population but may be important in some areas. For example, calf survival rates are lowest in the Cabinet-Salish population; however, the population appears to remain stable because of high adult female survival. Monitoring of moose vital rates as well as potential limiting factors (predation, disease, and nutrition) will continue for the remainder of this 10-year study.

Web site: We refer readers to our project website for additional information, reports, publications, photos and videos. The direct website for this moose study is: <http://fwp.mt.gov/fishAndWildlife/diseasesAndResearch/research/moose/populationsMonitoring/default.html>

Or alternatively, go to fwp.mt.gov. Click on the “Fish & Wildlife” tab at the top... then near the bottom right click on “Wildlife Research”... and follow links for “Moose”.

1.2. Evaluating genetic support for the Shiras subspecies of moose, *A. a. shirasi*

The taxonomic designation of subspecies is often brought to bear in the management and conservation of species, yet the definition and delineation of subspecies units have suffered from inconsistency across taxa and over time. We currently have a manuscript in revision at a peer-reviewed journal that applies a broader lens to the genetic structure of moose in the West. Specifically, we are applying contemporary guidelines of subspecies delineation according to mitochondrial genomics to evaluate the southernmost subspecies of moose, *Alces alces shirasi*. We sequenced the complete mitochondrial genome ($N=60$) as well as 13 nuclear microsatellites ($N=253$) from moose across western North America to evaluate the genetic distinction of moose within the putative range of the *A. a. shirasi* subspecies. Fossil records were consulted to supply phylogeographic context to genetic structure. Results from this study will become available upon acceptance of the manuscript, expected FY20.


Objective #2: Monitor moose vital rates and potential limiting factors

2.1. Animal capture and handling


In January of 2019 we worked with a contracted helicopter capture company (Quicksilver Air) and local landowners to conduct captures and increase the sample of monitored moose. A total of 24 adult females were captured in 3 study areas in 2018, with the goal of maintaining 30 collared animals in each area. Moose were fit with GPS radio-collars (Lotek LifeCycle and Vectronic Survey Globalstar). During 2013–19, we have conducted a total of 181 captures of 162 individual adult female moose, and as of August 1, 2019, 81 are currently being monitored (Table 1, Figure 2). A target sample size of 30 individuals/study area is sought to achieve moderate precision in annual survival estimates, while minimizing capture and monitoring costs.

Table 1. Captures of radio-marked adult female moose by study area and year, excluding 6 capture-related mortalities, and the number of adult females being monitored as of August, 2019.


	Study Area			Total
	Cabinet-Salish	Big Hole Valley	Rocky Mtn Front	
2013 captures	11	12	11	34
2014 captures	7	20	8	35
2015 captures	13	6	7	26
2016 captures	0	4	6	10
2017 captures	10	7	9	26
2018 captures	7	8	11	26
2019 captures	8	6	10	24
Total captures	56	63	62	181
Moose currently on-air	24	28	29	81



Cabinet-Salish Mountains



Big Hole Valley



Rocky Mountain Front




Figure 2. Moose winter capture locations during 2013–2019 across 3 study areas in Montana.

2.2. Monitoring vital rates

2.2.1. Adult female survival.— Our study of adult female survival to date includes 141 radio-collared adult female moose and 466 animal-years of monitoring, with a staggered-entry design of individuals entering into the study across 6 winter capture seasons (*see* 2.1 Animal capture and handling). Animals have been deployed with both VHF ($N=76$) and GPS ($N=86$) collars. We estimated Kaplan-Meier annual survival rates for each study area during each biological year as well as across the 5 biological years pooled together in a recurrent-time format.

Pooled annual survival estimates across the entire monitoring period for each study area were 0.891 (SE=0.024, 95% CI=[0.84,0.94]) in the Cabinet-Salish, 0.864 (SE=0.027, 95% CI=[0.81,0.92]) in the Big Hole Valley, and 0.890 (SE=0.025, 95% CI=[0.84,0.94]) on the Rocky Mountain Front (Figure 3). In comparison to these 5-year averages, survival during the 2018-19 biological year was low in the Cabinet-Salish (0.75), higher in the Big Hole Valley (0.93), and low on the Rocky Mountain Front (0.85). While differences among study areas appeared more pronounced during the early years of this study, the mean estimates in each area have gradually grown closer to one another as we continue to accumulate data. These estimates do not account for differences in age distribution of our collared sample, which we will address in more detail upon completion of the study (Prichard et al. 2012).




Figure 3. *Kaplan-Meier estimates of annual adult female survival within each study, where bolded lines are pooled estimates across 6 biological years for each study area and thin lines are annual estimates for each study area and year, Montana, 2013–2019.*

During 6 biological years of monitoring, we have documented 58 mortalities of collared adult moose across all study areas: 19 in the Cabinet-Fisher, 22 in the Big Hole Valley and 17 in the Rocky Mountain Front (Table 2, Figure 4). Ongoing research will attempt to better understand the causes and consequences of these mortalities.

Table 2. Numbers of mortalities by cause for radio-collared adult female moose documented during February 2013–June 2018, Montana.

Cause of Mortality	Study area		
	Cabinet-Fisher	Big Hole Valley	Rocky Mountain Front
Accident	1	0	0
Health-related (e.g., disease, malnutrition, ...)	6	20	8
Human (e.g., harvest, poaching, vehicle collision)	1	2	1
Predation, bear	0	0	1
Predation, mountain lion	1	0	0
Predation, wolf	6	0	2
Predation, wound infection	1	0	0
Unknown	3	0	5



Figure 4. Example mortality site of F347 in the Big Hole Valley study area, 10 February 2019. This individual was in poor nutritional condition, indicating a chronic health problem potentially related to a parasite load of ≥ 37 arterial worms in her carotid arteries. No other health-related abnormalities were detected, and she still had a calf with her at the time of death.

2.2.2 Calf survival.— We decompose calf survival into 2 components: 1) observed parturition rate – the proportion of pregnancies that result in a neonate calf-at-heel during spring; and 2) calf survival – the proportion of documented calves that survive through their first year of life.

Observed parturition rates: Following winter pregnancy testing, we use weekly aerial telemetry flights during 15 May – 15 July to estimate an “observed parturition” rate, representing the proportion of pregnant cows with neonate calves each spring. One limitation to this approach comes with the unknown proportion of the true number of calves born that die before we visually confirm them. Thus, our sample for subsequent study of calf survival is left truncated (Gilbert et al. 2014), and our Kaplan-Meier based estimates of calf survival should be considered as optimistic to the extent that they don’t account for mortality of calves prior to initial detection. Observed parturition rates have been higher in the Big Hole Valley (88%) and Rocky Mountain Front (92%), and lower in the Cabinet-Fisher (76%; Figure 5). These results are similar to those of other studies (e.g., Becker 2008) where parturition rates are lower than pregnancy rates due to presumed fetal losses throughout winter and/or death of neonatal calves prior to detection.

Calf survival: As a result of spring monitoring of neonate calves, we have documented 260 calves from 245 litters during 2013–2017. We then monitored the fates of these calves by visually locating them with their dams throughout their first year of life. Over the first 6 biological years (May 2013 – May 2019), pooled Kaplan-Meier survival estimates of calves-at-heel were 0.366 (SE=0.051, 95% CI=[0.28,0.48]) in the Cabinet-Fisher, 0.445 (SE=0.050, 95% CI=[0.36,0.55]) in the Big Hole Valley, and 0.526 (SE=0.049, 95% CI=[0.44, 0.63]) on the Rocky Mountain Front (Figure 5). Calf survival results mirror those of observed parturition, suggesting observed parturition rates are strongly influenced by mortality of neonates prior to detection, more so than fetal losses.




Figure 5. Observed parturition (proportion of pregnant cows with calves-at-heel during spring) and Kaplan-Meier estimates of annual calf survival for the first year of life within each study area, where bold lines are pooled estimates across 6 biological years and thin lines are annual estimates per year, Montana, 2013–2019.

2.2.3 Adult female fecundity.—Fecundity for moose is the product of age-specific pregnancy rates and litter size. We monitor pregnancy of animals during winter with laboratory analyses of both blood (serum PSPB levels; Huang et al. 2000) and scat (fecal progestagens; Berger et al. 1999, Murray et al. 2012). To estimate pregnancy in absence of handling animals each winter, we use fecal progestagens from samples collected via ground-tracking.

Pregnancy rates: Pooled across 3 study areas, 6 years (2013-2018), and 417 animal-years of monitoring, the average adult (ages ≥ 2.5) pregnancy rate was 82%, varying from 81–85% across study areas (Figure 6). Yearling (age 1.5) pregnancy rates appear to vary by region, with 0% pregnancy in both the Cabinet-Fisher and Big Hole Valley study areas compared to 45% yearling pregnancy on the Rocky Mountain Front; however, sample sizes for yearling pregnancy are small ($N = 2, 7,$ and 11 in the 3 areas, respectively).

Observed twinning rates: Moose are capable of giving birth to 1–3 calves, though litters are most commonly composed of either 1 or 2 calves (Van Ballenberghe and Ballard 2007). Twinning rates in North American populations can vary from 0 to 90% of births (Gasaway et al. 1992), with variation linked to nutritional condition (Franzmann and Schwartz 1985) and animal age (Ericsson et al. 2001). Twinning rates for Shiras moose are typically low (e.g., $<15\%$; Peek 1962, Schladweiler and Stevens 1973, Becker 2008). Thus far our observed twinning rates are 6% in the Cabinet-Fisher ($N=83$ litters), 1% in the Big Hole Valley ($N=105$ litters), and 16% in the Rocky Mountain Front study areas ($N=95$ litters; Figure 6).




Figure 6. Estimated adult (age ≥ 2.5) pregnancy rates, yearling (aged 1.5) pregnancy rates, observed twinning rates, and net observed fecundity of calves per adult female in 3 study areas of Montana during 6 biological years, 2013–2018.

2.2.4. Population growth rates. The overall status of a population may be best characterized by the annual growth rate. This parameter can be estimated by inserting key vital rates into mathematical models, most importantly the annual survival of adult females and the per capita number of calves born and surviving their first year. We estimated recruitment per cow as the integrated product of pregnancy rates, parturition rates, litter size, and calf survival rates. We then estimated annual population growth rates, following DeCesare et al. (2012), for each study population across the first 6 biological years, 2013–2019 (Figure 7).

Given the high elasticity of adult female survival in long-lived, iteroparous species (Eberhardt 2002), adult female survival is the most important vital rate for determining population growth rates. High adult survival, on average, in the Cabinet-Fisher translated to a mean population growth rate of 1.01, or an 1% increase per year, despite consistently seeing the lowest calf survival of all 3 areas. The Rocky Mountain Front moose have seen very high survival rates of both adults and calves as well as high fecundity of adults, resulting in an estimated annual growth rate of 1.12. While vital rates in the Big Hole Valley population were indicative of a declining population for several years, higher adult survival in recent years has increased the overall average to show a stable to increasing population growth rate (1.02) for the first time of the study.




Figure 7. Contour plot showing the estimated mean annual population growth rates (λ , represented as contour lines) resulting from two-dimensional combinations of adult female survival and spring recruitment of calves (integrating rates of pregnancy, parturition, litter size, and calf survival through the first year). Smaller dots show annual rates for each of 6 biological years, and larger dots and error bars show the pooled averages and standard errors, 2013–2019. Growth rates above the bold line (where $\lambda = 1$) indicate a growing population, growth rates below $\lambda = 1$ indicate declining populations. Results are preliminary and subject to change.

2.3. Monitoring nutritional condition, antler spreads, and rutting behavior with the voluntary help of moose hunters

2.3.1. Hunter-based sampling of nutritional condition.

Nutritional condition of ungulates can impact both survival (Roffe et al. 2001, Bender et al. 2008) and fecundity (Testa and Adams 1998, Keech et al. 2000, Testa 2004), and generally provides an indication of the extent to which habitat condition and density dependent effects drive ungulate dynamics (Franzmann and Schwartz 1985, Bertram and Vivion 2002). Rump fat thickness has a strong linear relationship with total body fat in moose (Stephenson et al. 1998). In addition to measuring rump fat among all captured adult females, since 2013 we have asked hunters to measure rump fat of harvested moose.

Moose hunters measured rump fat by marking a toothpick within provided sampling kits for 393 bull and 47 cow moose. Before comparing fat measurements across regions of Montana, we first assessed the relationship between the date each moose was harvested and its respective fat levels, as bull moose are known to lose fat with high energy expenditure during the rutting season (Cederlund et al. 1989). While there was much variation, we found a significant and consistent loss in rump fat depth among bull moose during each of the 5 years ($P < 0.001$), whereas fat among cows did not change with day of season ($P = 0.68$; Figure 8).




Figure 8. Depth of rump fat declined consistently among harvested bull moose according to the date of harvest during the past 6 hunting seasons (see 6 black trend-lines), whereas average fat depths among cow moose did not significantly change (red trend-line) during the hunting season, Montana, 2013–2018.

After assessing how average fat levels changed during the season, we compared observed measurements of fat for each moose to the average expected amount of fat following the trend lines in Figure 8. We then estimated the residuals between observed and predicted values, where a positive value suggested an animal with more fat than expected given the date of harvest, and a negative value an animal with less fat than expected. We compared these residual values among all MFWP regions and found no evidence for statistical differences in the nutritional conditions of bull moose among regions (Figure 9). However, we did find evidence of a difference in the rate at which bulls deplete their fat stores, according to their size (Figure 10). We also found evidence that moose increase in antler spread size up until about the age of 6 years old, at which point antler spreads generally are not affected by age (Figure 11).




Figure 9. Average residual values comparing the thickness of rump fat in hunter-killed moose among regions while controlling for the date of harvest. These data were collected by hunters by marking a toothpick (inset photo) included in sampling kits mailed to all license-holders, Montana, 2013–2018.




Figure 10. We also continue to find evidence from hunter-collected measurements of both rump fat and antler spread that larger bulls generally start the rutting season with more fat, but deplete their fat stores at a faster rate than smaller bulls, ending the rut in poorer nutritional condition, Montana, 2013–2018.



Figure 11. Hunter self-reported, unofficial, antler spreads (inches) show a gradual increase in the average spread across all moose statewide until about the age of 6, at which point average antler spread holds steady with age.

2.3.2. Hunter-based monitoring of the rut

For the lucky few (1.2% of applicants in 2018) who draw a moose license each year, one of the first considerations in hunt planning is the timing of the rut for moose in Montana. Mean breeding dates for moose in other studies have included October 5–10 in British Columbia, September 29 in Manitoba, and October 5 in Alaska (Schwartz 2007). During 2016–18, we asked moose hunters to mark on a calendar which days they hunted, and which days they observed rutting activity by moose (e.g., calling, sparring, wallowing). We received samples and/or information from an average of 157 hunters each year, including the recording of 3,632 hunter-days and 555 observations of rutting activity. Hunter-days decreased gradually throughout the season, with recurrent weekly spikes of hunting activity during weekends (Figure 12). To the contrary, the proportion of hunters observing rutting activity increased until the first week of October during both years, after which it declined through the middle of October (Figure 12). These observations are in accordance with our estimates of peak breeding based on estimated average parturition dates for radio-collared cows (May 23rd) and a 231-day gestation (Schwartz & Hundertmark 1993).





Figure 12. Hunter-days recorded from voluntary return of data cards and proportion of hunters observing moose rutting activity (using a 5-day moving average) throughout the hunting season, 2016-18, Montana.

2.4. Multi-species predator occupancy


Predation is among the hypothesized factors potentially limiting moose vital rates in Montana, and the extent to which predation limits moose populations is of widespread interest. Past research has found predation by grizzly bears, black bears and wolves could have potentially significant effects on moose populations, under some circumstances (Messier and Crête 1985, Larsen et al. 1989, Ballard et al. 1990). In addition, mountain lions are known to predate on moose and even coyotes may take calves (Ross and Jalkotzy 1996, Bartnick et al. 2013, Benson and Patterson 2013). Given the potential role of these carnivores in moose population dynamics, and perhaps more importantly the effects of the predator guild as a whole (Sih et al. 1998, Griffin et al. 2011, Keech et al. 2011), we are assessing the relationship between predator densities and moose vital rates. Of particular interest is the effects of predation effects on moose calf survival, which appears to differ between study areas (see pg.8). However, obtaining information on predator populations and their potential to effect moose survival is challenging because these species are cryptic and occur at low densities. Camera trapping is a promising means of obtaining estimates of occupancy and relative density for multiple species simultaneously in a manner that is non-invasive and cost-effective (Rovero & Marshal 2009, Brodie et al. 2014, Steenweg et al. 2016). This has led to rapid expansion of camera trapping for wildlife research and management, and concomitant statistical models to exploit camera data (Meek & Fleming 2014, Burton et al. 2015, Moeller et al. 2018).

Since September 2015 we have continuously operated remote camera grids on 3 moose field study areas to evaluate the ubiquity of predators, and the relationship between predator populations and moose vital rates. Remote cameras are deployed year-round in randomly selected cells within the trapping grid (Figure 13). Within the selected cell, unbaited camera sets are established on trails, closed roads and other travel routes to maximize detection of multiple carnivore species. Local landowners and managers have played an important role in the successful implementation of this research component. Along with providing access to areas, landowners and managers have contributed their knowledge and participation in field work to successfully establish and maintain camera sets. This work continues to foster new and existing relationships with landowners and managers in these areas. As of August 2019, we have deployed remote cameras at 130 sites (43 sites Cabinet-Salish study area; 49 Rocky Mountain Front study area; 38 Big Hole Valley study area) 85 of which are currently active (30 Cabinet-Salish; 28 Rocky Mountain Front; 27 Big Hole Valley). To date we have retrieved and stored images spanning ~85,000 active camera trap-days (35,000 Cabinet-Salish; 25,000 Rocky Mountain Front; and 25,000 Big Hole Valley). Classification of images, including identifying species and enumerating group sizes, is ongoing (Figure 14).


Analytical methods for estimating abundance of unmarked species from observation data is an active area of research (MacKenzie et al. 2002, Royle 2004, Chandler et al. 2013). Accurate estimates of annual abundances of each predator species would be ideal; however, a precise index which detects spatial and temporal heterogeneity in predator abundance/activity would also be useful in assessing the potential influence of predators on moose vital rates (Parsons et al. 2017, Keim et al. 2019). Our camera trapping efforts were designed to use occupancy-based models to estimate probability of occurrence over the study area; as well as extensions of occupancy models which use counts to estimate abundance (Royle 2004, Brodie et al. 2014, Fiske & Chandler 2017). These efforts are also amenable to alternative approaches, such as random encounter and time-to-event models, that also estimate abundance while attempting to account for detection probability (Rowcliffe et al. 2008, Moeller 2017). Additional information to parametrize detection probabilities of grizzly bears and wolves may be incorporated into models based on ongoing GPS telemetry monitoring by collaborators within MT FWP and the U.S. Fish and Wildlife Service.



Region 1: Cabinet-Fisher



Region 3: Big Hole Valley



Region 4: Rocky Mtn. Front




Figure 13. Sampling grids (2 x 2 miles) for deployment of remote cameras for monitoring multi-species predator occupancy across areas occupied by moose.




Figure 14. Example photos from remote camera-traps set within seasonal ranges of each moose study area to monitor multi-species occupancy of carnivores 2015–2019, Montana.

Deliverables

Below we list project deliverables (publications, reports, presentations, media communications, and value-added collaborations) stemming from this moose research project, during FYs 13–19 (July 2012–June 2019). In addition to those communications listed below, are frequent discussions with moose hunters statewide. Copies of reports and publications are available on the moose study’s website (note: the web address is case-sensitive):

<http://fwp.mt.gov/fishAndWildlife/diseasesAndResearch/research/moose/populationsMonitoring>

1. Annual Reports:

2013, 2014, 2015, 2016, 2017, 2018, 2019. DeCesare, N. J., and J. R. Newby. *Vital rates, limiting factors and monitoring methods for moose in Montana*. Annual reports, Federal Aid in Wildlife Restoration Grant W-157-R-1 through R-7.

2. Peer-reviewed Publications

Burkholder, B. O., N. J. DeCesare, R. A. Garrott, and S. J. Boccadori. 2017. *Heterogeneity and power to detect trends in moose browsing of willow communities*. *Alces* 53:23–39.

DeCesare, N. J., T. D. Smucker, R. A. Garrott, and J. A. Gude. 2014. *Moose status and management in Montana*. *Alces* 50:31–51.

DeCesare, N. J., J. R. Newby, V. Boccadori, T. Chilton-Radandt, T. Thier, D. Waltee, K. Podruzny, and J. A. Gude. 2016. *Calibrating minimum counts and catch per unit effort as indices of moose population trend*. *Wildlife Society Bulletin* 40:537–547.

Nadeau, M. S., N. J. DeCesare, D. G. Brimeyer, E. J. Bergman, R. B. Harris, K. R. Hersey, K. K. Huebner, P. E. Matthews, and T. P. Thomas. 2017. *Status and trends of moose populations and hunting opportunity in the western United States*. *Alces* 53:99–112.

Ruprecht, J. S., K. R. Hersey, K. Hafen, K. L. Monteith, N. J. DeCesare, M. J. Kauffman, and D. R. MacNulty. 2016. *Reproduction in moose at their southern range limit*. *Journal of Mammalogy* 97:1355–1365.

3. Other Publications

DeCesare, N. J. 2013. *Research: Understanding the factors behind both growing and shrinking Shiras moose populations in the West*. *The Pope and Young Ethic* 41(2):58–59.

DeCesare, N. J. 2014. *Conservation Project Spotlight: What and where are Shiras moose?* *The Pope and Young Ethic* 42(4):26–27.

4. Professional Conference Presentations

DeCesare, N. J., J. Newby, V. Boccadori, T. Chilton-Radant, T. Their, D. Waltee, K. Podruzny, and J. Gude. 2015. *Calibrating indices of moose population trend in Montana*. North American Moose Conference and Workshop, Granby, Colorado.

Nadeau, S., E. Bergman, N. DeCesare, R. Harris, K. Hersey, P. Mathews, J. Smith, T. Thomas, and D. Brimeyer. 2015. *Status of moose in the northwest United States*. North American Moose Conference and Workshop, Granby, Colorado.

DeCesare, N. J., J. R. Newby, and J. M. Ramsey. 2015. *A review of parasites and diseases impacting moose in North America*. Montana Chapter of the Wildlife Society. Annual Meeting, Helena, Montana.

DeCesare, N. J., J. Newby, K. Podruzny, K. Wash, and J. Gude. 2016. *Occupancy modeling of hunter sightings for monitoring moose in Montana*. North American Moose Conference and Workshop, Brandon, Manitoba.

Newby, J. R., N. J. DeCesare, and J. A Gude. 2016. *Assessing age structure, winter ticks, and nutritional condition as potential drivers of fecundity in Montana moose*. Montana Chapter of the Wildlife Society. Annual Meeting, Missoula, Montana.

Newby, J. R., N. J. DeCesare, and J. A Gude. 2016. *Assessing age structure, winter ticks, and nutritional condition as potential drivers of fecundity in Montana moose*. North American Moose Conference and Workshop, Brandon, Manitoba.

DeCesare, N. J., J. Newby, K. Podruzny, K. Wash, and J. Gude. 2017. *Occupancy modeling of hunter sightings for monitoring moose in Montana*. Montana Chapter of the Wildlife Society. Annual Meeting, Helena, Montana.

DeCesare, N. J., and J. R. Newby. 2018. *Moose population dynamics in Montana: results from the halfway point of a 10-year study*. Montana Chapter of the Wildlife Society. Annual Meeting, Butte, Montana.

Oyster, J. H., N. J. DeCesare, et al. 2018. *An update on Elaeophora schneideri in western North American moose*. North American Moose Conference and Workshop, Spokane, Washington.

DeCesare, N. J., and J. R. Newby. 2018. *Moose population dynamics in Montana*. North American Moose Conference and Workshop, Spokane, Washington.

DeCesare, N. J., et al. 2019. *Phylogeography of a range edge subspecies: is there such thing as Shiras moose?* Montana Chapter of the Wildlife Society. Annual Meeting, Helena, Montana.

5. Public and/or Workshop Presentations

FY	Organization (<i>Speaker</i>)	Location
2013	Helena Hunters and Anglers Association (<i>DeCesare</i>)	Helena, MT
	Marias River Livestock Association (<i>DeCesare</i>)	Whitlash, MT
	Plum Creek Timber Company, Staff meeting (<i>DeCesare</i>)	Libby, MT
	Sun River Working Group (<i>DeCesare</i>)	Augusta, MT

2014	Big Hole Watershed Committee (<i>DeCesare</i>)	Divide, MT
	Flathead Wildlife Incorporated (<i>DeCesare</i>)	Kalispell, MT
	MFWP R1, Regional Citizens Advisory Council (<i>Newby</i>)	Kalispell, MT
	MFWP R1, Biologists' Meeting (<i>Newby</i>)	Kalispell, MT
	MFWP R1, Bow Hunter Education Workshop	Kalispell, MT
	MFWP R2, Regional Meeting (<i>DeCesare</i>)	Missoula, MT
	MFWP, Wildlife Division Meeting (<i>DeCesare</i>)	Fairmont, MT
	Plum Creek Timber Annual Contractors Meeting (<i>DeCesare</i>)	Kalispell, MT
	Rocky Mountain Front Land Managers Forum (<i>DeCesare</i>)	Choteau, MT
	Swan Ecosystem Center Campfire Program (<i>Newby</i>)	Holland Lake, MT
	WCS Community Speaker Series (<i>Newby</i>)	Laurin, MT
2015	Big Hole Watershed Committee (<i>Boccadori</i>)	Divide, MT
	Flathead Chapter of Society of American Foresters (<i>Newby</i>)	Kalispell, MT
	Libby Chapter of Society of American Foresters (<i>Newby</i>)	Libby, MT
	MFWP R1, Regional Citizens Advisory Council (<i>Newby</i>)	Kalispell, MT
	MFWP R2, Bow Hunter Education Workshop (<i>DeCesare</i>)	Lolo, MT
	MFWP R2, Regional Citizens Advisory Council (<i>DeCesare</i>)	Missoula, MT
	Rocky Mountain Front Land Managers Forum (<i>Newby</i>)	Choteau, MT
	Sanders County Commission Meeting (<i>DeCesare</i>)	Thompson Falls, MT
	Sheridan Wildlife Speaker Series (<i>DeCesare</i>)	Sheridan, MT
	Univ. Montana Guest Lecture – WILD105 (<i>DeCesare</i>)	Missoula, MT
2016	Confederated Salish & Kootenai Tribe, Nat Res Commission (<i>Newby</i>)	Marion, MT
	Ducks Unlimited State Convention (<i>Newby</i>)	Lewistown, MT
	Helena Hunters and Anglers Association (<i>DeCesare</i>)	Helena, MT
	MFWP R1 Law Enforcement Annual Meeting (<i>Newby</i>)	Kalispell, MT
	Montana State University, Ecology Seminar Series (<i>DeCesare</i>)	Bozeman, MT
	Ravalli County Fish and Wildlife Association (<i>DeCesare</i>)	Hamilton, MT
	Univ. Montana Guest Lecture – WILD480 (<i>DeCesare</i>)	Missoula, MT
	Upper Sun River Wildlife Team Meeting (<i>DeCesare</i>)	August, MT
2017	Big Hole Watershed Committee (<i>Boccadori</i>)	Divide, MT
	Mountain Bluebird Trails Conference (<i>DeCesare</i>)	Dillon, MT
	Swan Valley Connections Speaker Series (<i>DeCesare</i>)	Condon, MT
	University of Montana, STEAMfest (<i>DeCesare</i>)	Missoula, MT
	Univ. Montana Guest Lectures – WILD180, WILD480 (<i>DeCesare</i>)	Missoula, MT
	WCS Community Speaker Series (<i>DeCesare</i>)	Dillon, MT
	Flathead Valley Lions Club (<i>Newby</i>)	Kalispell, MT
	Flathead Wildlife Incorporated (<i>Newby</i>)	Kalispell, MT
	North Fork Inter-local (<i>Anderson</i>)	Polebridge, MT
2018	Bitterroot College (<i>DeCesare</i>)	Hamilton, MT
	Clearwater Resource Council (<i>DeCesare</i>)	Seeley Lake, MT
	MFWP R1, Regional Citizens Advisory Council (<i>Newby</i>)	Kalispell, MT
	Montana Forest Landowner Conference (<i>DeCesare</i>)	Helena, MT
	Montana Audubon Chapter (<i>Newby</i>)	Polson, MT
	Science on Tap (<i>Newby</i>)	Bigfork, MT
2019	MFWP HQ, Brown Bag Seminar (<i>DeCesare</i>)	Helena, MT
	MFWP Wildlife Manager Meeting (<i>DeCesare</i>)	Helena, MT
	Hellgate Hunters and Anglers (<i>DeCesare</i>)	Missoula, MT
	Rocky Mountain Front Land Managers Forum (<i>Newby</i>)	Choteau, MT

Upper Sun Wildlife Team (<i>DeCesare</i>)	Fairfield, MT
Univ. Montana Guest Lectures – WILD240 (<i>DeCesare</i>)	Missoula, MT
Idaho Fish & Game/MFWP Joint Meeting (<i>Newby</i>)	De Borgia, MT

6. Media Communications

FY	Organization (Location)	Topic	Media
2013	Bozeman Chronicle (MT)	Moose research	Newspaper
	Liberty County Times (MT)	Moose research	Newspaper
	MFWP Outdoor Report (MT)	Moose research	Television
2014	Carbon County News (MT)	Moose research	Newspaper
	Flathead Beacon (MT)	Moose research	Newspaper
	Helena Independent Record (MT)	Moose research	Newspaper
	High Country News, blog	Moose research	Blog
	KPAX (MT)	Moose-human conflict	Television
	MFWP Outdoor Report	Moose research	Television
	Missoulian (MT)	Urban moose	Newspaper
	The Monocle Daily (London, UK)	Moose research	Radio
	Nature Conservancy Magazine (VA)	Moose research	Magazine
	New York Times (NY)	Moose research	Newspaper
	NWF Teleconference (MT)	Climate change	Newspaper
	Radio New Zealand (New Zealand)	Moose research	Radio
	Summit Daily (CO)	Moose research	Newspaper
	UM Science Source (MT)	Moose research	Newspaper
2015	KOFI (MT)	Moose research	Radio
	MFWP Outdoor Report (MT)	Moose research	Television
	Western News (MT)	Moose research	Newspaper
2016	Missoulian (MT)	Climate & moose	Newspaper
	Bozeman Daily Chronicle (MT)	Climate & moose	Newspaper
	Montana Standard (MT)	Climate & moose	Newspaper
	Billings Gazette (MT)	Climate & moose	Newspaper
	Daily Interlake (MT)	Moose research	Newspaper
	Ravalli Republic (MT)	Moose research	Newspaper
	Montana Public Radio (MT)	Moose research	Radio
	Montana Public Radio – Field Notes (MT)	Moose taxonomy	Radio
	Post Rider (MT)	Moose research	Newsletter
	KAJ18 (MT)	Moose research	Television
	2017	Dillon Tribune (MT)	Moose research
Billings Gazette (MT)		Moose research	Newspaper
Missoulian (MT)		Moose research	Newspaper
Great Falls Tribune (MT)		Moose research	Newspaper
Weather Network (Canada)		Moose sightings	Website
The Nature Conservancy Magazine (VA)		Wildlife tracking	Magazine
2018	Hungry Horse News (MT)	Moose research	Newspaper
	Missoulian (MT)	Moose research	Newspaper
2019	Missoulian (MT)	Moose hunting	Newspaper

7. Other Project-related Collaborations

Partners	Title	Activities during FY18
Rick Gerhold & Caroline Grunenwald, University of Tennessee	Development of a serological assay for <i>Elaeophora schneideri</i> detection and surveillance in cervids	*Labwork is ongoing *Providing MT blood samples and worm samples for lab work
Biologists from western states and provinces (AB, BC, CO, ID, MT, OR, SK, UT, WA, WY)	Assessing range-wide genetic differentiation and spatial distribution of a moose subspecies, <i>Alces alces shirasi</i>	*Manuscript 2 nd revision in review at Journal of Mammalogy
Biologists from western states (CO, ID, MT, OR, UT, WA, WY)	Summarize status and management of western states moose.	*Completed, manuscript published, 2017.
Ky Koitzsch, K2 Consulting, LLC	Estimating population demographics of moose in northern Yellowstone National Park using non-invasive methods	*Field work and analyses completed *Final report in development
Jason Ferrante & Margaret Hunter, USGS – Gainseville, FL	Genetic approaches to understanding moose health	*Analyses ongoing

Acknowledgements

Private landowners & hunters: We are particularly thankful to The Nature Conservancy, Stoltze Land & Lumber, Weyerhaeuser and the many private landowners and area residents who graciously allowed us to conduct captures and ground monitoring on their properties. They also have provided much logistical support and local knowledge about moose distributions and local flora and fauna. We are very grateful for the privilege to work on these properties and for all the help. We are also grateful to the many moose hunters who have given their time and effort to submit samples and information to us as part of their moose hunting experience.

This project is a large collaboration among many FWP biologists. These include but are not limited to Justin Gude, Jennifer Ramsey, Neil Anderson, Keri Carson, Kevin Podruzny, Keri Wash, Jim Williams, Howard Burt, Graham Taylor, Tonya Chilton-Radant, Kent Laudon, John Vore, Vanna Boccadori, Brent Lonner, Gary Olson, Jessy Coltrane, and Ryan Rauscher. Undoubtedly this list should be larger to fully incorporate the many biologists and other personnel who have assisted with coordination of hunter sample collection, harvest statistics,

opportunistic sampling of other moose throughout the state. We also acknowledge a great deal of help from other cooperating biologists and agency personnel including Nathan Birkeland, Dave Hanna, Dan Carney, Lorin Hicks, Allison Kolbe, Jenna Roose, Michael Munoz, and Eric Eneboe, Alissa Anderson and Kaitlyn Farrar. Thanks also to Caryn Dearing, Adam Brooks, Kari Shinn, Quentin Kujala, and others at FWP-HQ for their help with project administration.

Many thanks go to the pilots who have safely conducted capture and telemetry work, including Mike Campbell, Rob Cherot, Rick Geiger, Ken Justus, Blake Malo, Jim-Bob Pierce, Joe Rahn, Mark Stott, Rick Swisher, Guy Terrill, and Trever Throop.

Funding for this project during FYs 13–19 has been in part derived from FWP moose license auction sales, matched with USFWS Grants-in-Aid funds. We also thank the Safari Club International Foundation for 2 Conservation Grants to support this research during FY16–19. Thanks to Ducks Unlimited and the Rocky Mountain Elk Foundation for holding license auctions. Many thanks also go to Plum Creek Timber Company and Two Bear Aviation, which have supported the project with helicopter flights during capture and monitoring efforts.

Literature cited

- Ballard, W. B., S. D. Miller, and J. S. Whitman. 1990. Brown and black bear predation on moose in southcentral Alaska. *Alces* 26:1–8.
- Bartnick, T. D., T. R. Van Deelen, H. B. Quigley, and D. Craighead. 2013. Variation in cougar (*Puma concolor*) predation habits during wolf (*Canis lupus*) recovery in the southern Greater Yellowstone Ecosystem. *Canadian Journal of Zoology* 91:82–93.
- Becker, S. A. 2008. Habitat selection, condition, and survival of Shiras moose in northwest Wyoming. University of Wyoming.
- Bender, L. C., J. G. Cook, R. C. Cook, and P. B. Hall. 2008. Relations between nutritional condition and survival of North American elk *Cervus elaphus*. *Wildlife Biology* 14:70–80.
- Benson, J. F., and B. R. Patterson. 2013. Moose (*Alces alces*) predation by eastern coyotes (*Canis latrans*) and eastern coyote x eastern wolf (*Canis latrans* x *Canis lycaon*) hybrids. *Canadian Journal of Zoology* 91:837–841.
- Berger, J., J. W. Testa, T. Roffe, and S. L. Monfort. 1999. Conservation endocrinology: a noninvasive tool to understand relationships between carnivore colonization and ecological carrying capacity. *Conservation Biology* 13:980–989.
- Bertram, M. R., and M. T. Vivion. 2002. Moose mortality in eastern interior Alaska. *The Journal of wildlife management* 66:747–756.
- Boer, A. H. 1992. Fecundity of North American moose. *Alces* (Supplement) 1:1–10.
- Brodie, J. F., A. J. Giordano, B. Dickson, M. Hebblewhite, H. Bernard, J. Mohd-Azlan, J. Anderson, and L. Ambu. 2014. Evaluating Multispecies Landscape Connectivity in a Threatened Tropical Mammal Community. *Conservation Biology*.
- Burton, A. C., E. Neilson, D. Moreira, A. Ladle, R. Steenweg, J. T. Fisher, E. Bayne, and S. Boutin. 2015. REVIEW: Wildlife camera trapping: a review and recommendations for linking surveys to ecological processes. *Journal of Applied Ecology* 52:675–685.
- Cederlund, G. N., R. L. Bergström, and K. Danell. 1989. Seasonal variation in mandible marrow fat in moose. *The Journal of Wildlife Management*:587–592.
- Chandler, R. B., J. A. Royle, and others. 2013. Spatially explicit models for inference about density in unmarked or partially marked populations. *The Annals of Applied Statistics* 7:936–954.
- DeCesare, N. J., M. Hebblewhite, M. Bradley, K. G. Smith, D. Hervieux, and L. Neufeld. 2012. Estimating ungulate recruitment and growth rates using age ratios. *Journal of Wildlife Management* 76:144–153.
- Eberhardt, L. L. 2002. A paradigm for population analysis of long-lived vertebrates. *Ecology* 83:2841–2854.
- Ericsson, G., K. Wallin, J. P. Ball, and M. Broberg. 2001. Age-related reproductive effort and senescence in

- free-ranging moose, *Alces alces*. *Ecology* 82:1613–1620.
- Franzmann, A. W., and C. C. Schwartz. 1985. Moose twinning rates: a possible population condition assessment. *The Journal of wildlife management* 49:394–396.
- Gasaway, W. C., R. D. Boertje, D. V. Grangaard, D. G. Kelleyhouse, R. O. Stephenson, and D. G. Larsen. 1992. The role of predation in limiting moose at low densities in Alaska and Yukon and implications for conservation. *Wildlife Monographs*:3–59.
- Gilbert, S.L., Lindberg, M.S., Hundertmark, K.J., Person, D.K., 2014. Dead before detection: addressing the effects of left truncation on survival estimation and ecological inference for neonates. *Methods in Ecology and Evolution* 5, 992–1001.
- Griffin, K. A., M. Hebblewhite, H. S. Robinson, P. Zager, S. M. Barber-Meyer, D. Christianson, S. Creel, N. C. Harris, M. A. Hurley, D. H. Jackson, B. K. Johnson, W. L. Myers, J. D. Raithel, M. Schlegel, B. L. Smith, C. White, and P. J. White. 2011. Neonatal mortality of elk driven by climate, predator phenology and predator community composition. *Journal of Animal Ecology* 80:1246–1257.
- Hochachka, W. M., D. Fink, R. A. Hutchinson, D. Sheldon, W.-K. Wong, and S. Kelling. 2012. Data-intensive science applied to broad-scale citizen science. *Trends in Ecology & Evolution* 27:130–137.
- Keech, M. A., R. T. Bowyer, M. Jay, V. Hoef, R. D. Boertje, B. W. Dale, and T. R. Stephenson. 2000. Life-history consequences of maternal condition in Alaskan moose. *The Journal of Wildlife Management* 64:450–462.
- Keech, M. A., M. S. Lindberg, R. D. Boertje, P. Valkenburg, B. D. Taras, T. A. Boudreau, and K. B. Beckmen. 2011. Effects of predator treatments, individual traits, and environment on moose survival in Alaska. *Journal of Wildlife Management* 75:1361–1380.
- Keim, J.L., Lele, S.R., DeWitt, P.D., Fitzpatrick, J.J., Jenni, N.S., 2019. Estimating the intensity of use by interacting predators and prey using camera traps. *Journal of Animal Ecology* 88, 690–701.
- Kéry, M. 2018. Identifiability in *N*-mixture models: a large-scale screening test with bird data. *Ecology* 99:281–288.
- Larsen, D. G., D. A. Gauthier, and R. L. Markel. 1989. Causes and rate of moose mortality in the southwest Yukon. *Journal of Wildlife Management*:548–557.
- MacKenzie, D. I., J. D. Nichols, J. E. Hines, M. G. Knutson, and A. B. Franklin. 2003. Estimating site occupancy, colonization, and local extinction when a species is detected imperfectly. *Ecology* 84:2200–2207.
- MacKenzie, D. I., J. D. Nichols, G. B. Lachman, S. Droege, J. Andrew Royle, and C. A. Langtimm. 2002. Estimating site occupancy rates when detection probabilities are less than one. *Ecology* 83:2248–2255.
- Messier, F., and M. Crête. 1985. Moose-wolf dynamics and the natural regulation of moose populations. *Oecologia* 65:503–512.
- Meek, P., Fleming, P., 2014. *Camera Trapping: wildlife management and research*. CSIRO Publishing, Collingwood, Australia.
- Moeller, A.K., 2017. *New methods to estimate abundance from unmarked populations using remote camera trap data (Thesis)*. University of Montana, Missoula, Montana.
- Murray, D. L., K. F. Hussey, L. A. Finnegan, S. J. Lowe, G. N. Price, J. Benson, K. M. Loveless, K. R. Middel, K. Mills, D. Potter, A. Silver, M.-J. Fortin, B. R. Patterson, and P. J. Wilson. 2012. Assessment of the status and viability of a population of moose (*Alces alces*) at its southern range limit in Ontario.
- Parsons, A.W., T. Forrester, W.J. McShea, M.C. Baker-Whatton, J.J. Millspaugh, R. Kays. 2017. Do occupancy or detection rates from camera traps reflect deer density? *J Mammal* 98, 1547–1557. <https://doi.org/10.1093/jmammal/gyx128>
- Peek, J. M. 1962. Studies of moose in the Gravelly and Snowcrest Mountains, Montana. *The Journal of Wildlife Management* 26:360–365.
- Prichard, A. K., K. Joly, and J. Dau. 2012. Quantifying telemetry collar bias when age is unknown: a simulation study with a long-lived ungulate. *Journal of Wildlife Management* 76:1441–1449.
- Roffe, T. J., K. Coffin, and J. Berger. 2001. Survival and immobilizing moose with carfentanil and xylazine. *Wildlife Society Bulletin* 29:1140–1146.
- Ross, P. I., and M. G. Jalkotzy. 1996. Cougar predation on moose in southwestern Alberta. *Alces* 32:1–8.
- Rovero, F., Marshall, A.R., 2009. Camera trapping photographic rate as an index of density in forest ungulates. *J. Appl. Ecol.* 46, 1011–1017.

- Rowcliffe, J.M., Field, J., Turvey, S.T., Carbone, C., 2008. Estimating animal density using camera traps without the need for individual recognition. *Journal of Applied Ecology* 45, 1228–1236.
- Royle, J. A. 2004. N-Mixture Models for Estimating Population Size from Spatially Replicated Counts. *Biometrics* 60:108–115.
- Schladweiler, P., and D. R. Stevens. 1973. Reproduction of Shiras moose in Montana. *The Journal of Wildlife Management*:535–544.
- Schwartz, C. C. 2007. Reproduction, natality and growth. Pages 141–171 *Ecology and management of the North American moose*. 2nd edition. University Press of Colorado, Boulder, Colorado.
- Schwartz, C. C., and K. J. Hundertmark. 1993. Reproductive characteristics of Alaskan moose. *The Journal of Wildlife Management* 57:454-468.
- Sih, A., G. Englund, and D. Wooster. 1998. Emergent impacts of multiple predators on prey. *Trends in ecology & evolution* 13:350–355.
- Steenweg, R., Whittington, J., Hebblewhite, M., Forshner, A., Johnston, B., Petersen, D., Shepherd, B., Lukacs, P.M., 2016. Camera-based occupancy monitoring at large scales: power to detect trends in grizzly bears across the Canadian Rockies. *Biological Conservation* 201, 192–200.
- Stephenson, T. R., K. J. Hundertmark, C. C. Schwartz, and V. Van Ballenberghe. 1998. Predicting body fat and body mass in moose with ultrasonography. *Canadian Journal of Zoology* 76:717–722.
- van Strien, A. J., C. A. M. van Swaay, and T. Termaat. 2013. Opportunistic citizen science data of animal species produce reliable estimates of distribution trends if analysed with occupancy models. *Journal of Applied Ecology* 50:1450–1458.
- Testa, J. W. 2004. Population dynamics and life history trade-offs of moose (*Alces alces*) in south-central Alaska. *Ecology* 85:1439–1452.
- Testa, J. W., and G. P. Adams. 1998. Body condition and adjustments to reproductive effort in female moose (*Alces alces*). *Journal of Mammalogy* 79:1345–1354.
- Van Ballenberghe, V., and W. B. Ballard. 2007. Population dynamics. Pages 223–245 *Ecology and management of the North American moose*. 2nd edition. University Press of Colorado, Boulder, Colorado.